



EJP SOIL
European Joint Programme

Towards climate-smart sustainable management of agricultural soils

Deliverable 2.3

Synthesis on estimates of achievable soil carbon sequestration on agricultural land across Europe

Due date of deliverable: M12
Actual submission date: 29.01.2021



This project has received funding from the European Union's Horizon 2020 research and innovation programme under grant agreement N° 862695

GENERAL DATA

Grant Agreement: 862695

Project acronym: EJP SOIL

Project title: Towards climate-smart sustainable management of agricultural soils

Project website: www.ejpsoil.eu

Start date of the project: February 1st, 2020

Project duration: 60 months

Name of lead contractor: INRAE

Funding source: H2020-SFS-2018-2020 / H2020-SFS-2019-1

Type of action: European Joint Project COFUND

DELIVERABLE NUMBER:	2.3
DELIVERABLE TITLE:	Synthesis on estimates of achievable soil carbon sequestration on agricultural land across Europe
DELIVERABLE TYPE:	Report/ORDP/Website/Ethics
WORK PACKAGE N:	WP2
WORK PACKAGE TITLE:	A roadmap for Agricultural Soil Management in Europe
DELIVERABLE LEADER:	Agroscope and CRA-W
AUTHORS:	Leonor Rodrigues, Julia Fohrafellner, Briec Hardy, Bruno Huyghebaert, Jens Leifeld
CONTRIBUTORS OF DATA AND CO-AUTHORES:	Alberto Sanz Cobeña ,Alice Budai, Andis Lazdiņš, Arezoo Taghizadeh, Axel Don, Bartosz Adamczyk, Benjamin Gimeno, Benjamin Sánchez, Bo Stenberg, Claudia Di Bene, Corina Carranca, Dalia Feziene, Daniel Rasse, Daria Seitz, Dario Fornara, Eduardo Aguilera, Elena Rodriguez, Eloïse Mason, Erich Inselsbacher, Gabriela Barančíková, Grzegorz Siebielec, Heide Spiegel, Imants Kukuļs Jacek Niedźwiecki, Jan Peter Lesschen, Karin Kauer, Kestutis Armolaitis, Lilian OSullivan, Lenka Pavlu, Maarten De Boever, Nils Kauer, Peter Kuikman, Peter Laszlo, Raquel Mano, Raimonds Kasparinskis, Rok Mihelič, Sevinc Madenoglu, Sophie Cornu, Sophie Zechmeister-Boltenstern, Stephan Glatzel Sylvain Pellerin, Teresa Gómez de la Bárcena, Thalisa Slier, Thomas Kätterer , Martin A. Bolinder, Kerstin Berglund, Toth Gergely
DISSEMINATION LEVEL:	PU



ABSTRACT

This synthesis identifies the available knowledge of achievable carbon sequestration in mineral soils and GHGs mitigation in organic soils in agricultural land, including pasture/grassland across Europe. The inventory of past and current studies on carbon sequestration and GHGs mitigation measures in agricultural soils and the methodology used for the assessment were considered from 25 Member states (MS) across Europe. The stocktake shows that availability of datasets concerning soil carbon sequestration (SCS) is variable among Europe. While northern Europe and central Europe is relatively well studied, there is a lack of studies comprising parts of Southern, Southeaster and Western Europe. Further, it can be concluded that at present country based knowledge and engagement is still poor; very few countries have an idea on their national-wide achievable carbon sequestration potential. The presented national SCS potentials (MS n=13) do however point towards important contributions to mitigate climate change by covering considerable shares of national greenhouse gas emissions from the agricultural sector in the range of 1-28.5 %, underpinning the importance of further investigations. In contrast to mineral soils, effective mitigation measures for organic soils while maintaining industrial agricultural production are still in its infancy. Very few mitigation options exist to mitigate GHG emissions without compromising agricultural production. Most GHG mitigation practices reported by the MS involve the restoration of organic soils, which means a complete abandonment of land from any agricultural use. Only one contribution (NL) reports possible mitigation potentials, which are based on specific water management measures (water level fixation). Nevertheless, there is an increasing awareness of the need of mitigation measures reflected by the several ongoing research projects on peatland management.



Table of contents

ABSTRACT	3
Table of contents.....	4
List of Tables	5
List of Figures.....	5
List of acronyms and abbreviations.....	6
1. Executive summary	7
2. Introduction and goal of T2.4.3.....	8
3. Methodology and data source	9
4. Results	11
4.1 Data and data coverage	11
4.2 Mineral soils	11
4.2.1 Question 1 (Q1): Are there any quantitative estimations of the achievable carbon sequestration under different soil management in your country (finished or ongoing studies)?	15
4.2.2 Measures and land use Q1	21
4.2.3 Methods Q1:.....	21
4.2.4 Question 2 (Q2): Are there any estimated C sequestration rates for different management practices derived from field studies in your country (finished or ongoing studies)?.....	24
4.2.5 Measures Q2	24
4.2.6 Methods for quantifying changes in C stocks (Q2)	25
4.3 Organic soils	27
4.3.1 Question 3 (Q3): Are there any quantitative estimates for reducing GHG emissions (CO ₂ , CH ₄ , N ₂ O) from managed organic soils under different soil management in your country (finished or ongoing studies)?	28
4.3.2. Question 4: Are there any estimated GHG mitigation potentials for managed organic soils derived from field studies in your country?	29
4.4 Knowledge gaps reported by MS	31
4.3.1 Knowledge gaps in C-Sequestration in mineral soils.....	31
4.3.2 Knowledge gaps in organic soils.....	38
4.5 General knowledge gaps and hot topics.....	42
4.5.1 GHG mitigation measure of organic soils:.....	42
4.5.2 General methodological challenges:	42
4.5.3 Socio-economic factors:	42
4.5.4 SCS under climate change	43
4.5.5 Trade off C-Sequestration vs other GHG:.....	43
4.5.6 Measures and alternatives:.....	44
4.5.7 Soil (bio-) chemical properties.....	44
4.5.8 Carbon use efficiency in differently managed systems.....	45
6 Conclusions.....	45
7 Limitations of the synthesis.....	46
References.....	46
Annex.....	55
A List of references mineral soils.....	55
B List of references organic soils	68
C List of finished and ongoing projects	72
D List of experts and contacts Task T 2.4.3	79
E Questionnaire for MS	81



List of Tables

Table 1 Agricultural measures to improve SCS in mineral soils mentioned in the stocktake.....	14
Table 2 Achievable domestic soil carbon sequestration (SCS) and reduction potential of the total emissions from the agricultural sector (EA) reported by the MS.....	18
Table 3 Counts of knowledge gap mentions per group and MS.....	31
Table 4 Counts of knowledge gaps per European region	32
Table 5 Number of knowledge gaps for organic soils and member states per group	39
Table 6 Knowledge gaps for organic soils per main European region	39

List of Figures

Figure 1 The Environmental Stratification of Europe.....	10
Figure 2 Type of Literature considering SCS and distribution from 1990-2021.....	12
Figure 3 Data density (Studies per 10'000km ²) for each environmental zone (EZ) across Europe.....	13
Figure 4 Share of potentials to the annual total emissions from the agricultural sector for each country and share of measures involved.	17
Figure 5 Share of measures mentioned for national estimates of SCS. Question 1 (Qs1).....	21
Figure 6 Share of measures mentioned for national estimates of SCS. Question 2 (Qs2).....	24
Figure 7 Rates derived from provided studies	26
Figure 8 Counts for time periods over which SCS was estimated.	27
Figure 9 Knowledge gap groups for mineral soils and the corresponding EJP SOIL challenges.	38
Figure 10 Knowledge gap groups for organic soils and the corresponding EJP SOIL challenges.....	41



List of acronyms and abbreviations

AGS	Agroscope
CRA-W	Centre Wallon de Recherches Agronomiques
CUE	carbon use efficiency
EA	Emissions from the agricultural sector
EZ	Environmental Zones: <ul style="list-style-type: none">• ALN Alpine North• ALS Alpine South• ANA Anatolian• ATC Atlantic Central• ATN Atlantic North• BOR Boral• CON Continental• LUS Lusitanian• NEM Nemoral• MDM Mediterranean mountains• MDN Mediterranean North• MDS Mediterranean South• PAN Panonian
GHG	Greenhouse gas
GL	Grey Literature
IPCC	Intergovernmental Panel on Climate Change
LTE	Long-term Experiment
MS	Member states
NIR	National Inventory Report
PSL	Published Scientific Literature
QS	Question
SCS	Soil Carbon Sequestration
SOC	Soil organic Carbon



1. Executive summary

This document summarizes the stocktaking activities for Europe led by Agroscope (AGS) and Centre Wallon de Recherches Agronomiques (CRA-W) in collaboration with 23 of the EJP SOIL consortium members. The goal of T2.4.3. is stocktaking and synthesis on available knowledge of achievable carbon sequestration in mineral soils and greenhouse gas (GHG) mitigation in organic soils in agricultural land, including pasture/grassland across Europe. All kind of information such as data, reports, scientific papers, including commercial and grey literature for each participating country were collected. The inventory of past and current studies on soil carbon sequestration (SCS) and GHGs mitigation measures in agricultural soils and the methodology used for the assessment were also considered. All participating MS (n=25) returned their country specific information. The stocktake shows that availability of datasets concerning SCS is variable among Europe. While northern Europe and central Europe is relatively well studied, there is a lack of studies comprising parts of Southern, Southeaster and Western Europe. There is however a clear increase in interest for topics covering SCS for climate change mitigation in the last two decades. A total of 114 ongoing projects have been reported dealing with SCS and GHG mitigation measures. However, at present country based knowledge and engagement is still poor. The stocktake shows only half of the participating MS (BE, CH, DE, DK, ES, FR, IR, IT, NL, NO, PO, SW, UK) have an idea on their national-wide achievable carbon sequestration potential. Information provided is mostly based on rough estimates without consideration of technical and socio-economic feasibilities (i.e. availability and suitability of land, costs or profits for farmers and the willingness of implementation by farmers), which are key for sustainable implementation. Climate change is usually not included, although it is known that it will have an important impact on carbon stocks. The presented national SCS potentials (n=13) do however point towards important contributions to mitigate climate change by covering considerable shares of national greenhouse gas emissions from the agricultural sector in the range of 1-28.5 %, underpinning the importance of further investigations.

In contrast to mineral soils, effective mitigation measures for organic soils while maintaining agricultural production are hardly known. Very few options exist to mitigate GHG emissions without compromising intensive agricultural production. Most GHG mitigation practices reported by the MS involve the restoration of organic soils, which implies abandonment of land from most agricultural uses. Only one contribution (NL) reports possible mitigation potentials, which are based on specific water management measures. Nevertheless, there is an increasing awareness of the need of mitigation measures reflected by the several ongoing research projects on peatland management.



2. Introduction and goal of T2.4.3

In the context of climate change and rising atmospheric CO₂ concentrations, the role of soils in the global carbon cycle has been increasingly acknowledged. Depending on land use and management, soils can act as both a source and a sink for CO₂ (Lal, 2004). Soils are the largest terrestrial pool of organic carbon with global soil organic carbon (SOC) stocks of 863 Pg C, 1,824 Pg C, and 3,012 Pg C in the upper 0.3 m, 1 m, and 2 m respectively (Sanderman et al., 2017). Several long-term soil-monitoring networks and data from long-term experiments (LTEs), in a considerable number of European countries, show that soil carbon in many agricultural soils is decreasing (Bellamy et al., 2005; Heikkinen et al., 2013; Keel et al., 2019; Sleutel et al., 2007; Taghizadeh-Toosi et al., 2014). However there is a general agreement that selected agricultural practices could have a significant positive impact on soil carbon sequestration (SCS), with further benefits for soil fertility and ecosystem services (e.g. water infiltration and -holding capacity, provision of food and ecological resilience). Hence, since the last two decades, agricultural practices have been increasingly endorsed as an option to remove CO₂ from the atmosphere to mitigate climate change and a growing number of literature reviews show that specific agricultural measures can increase SCS (Lal, 2004; Smith et al., 2007, 2005). However, an increase in soil organic matter does not necessarily imply carbon (C) to be sequestered from the atmosphere. Within the EJP SOIL SCS is defined as “the process of transferring CO₂ from the atmosphere into the soil of a land unit, through plants, plant residues and other organic solids which are stored or retained in the unit as part of the soil organic matter for a given time period” (Olson et al., 2014).

LTEs and modelling are general approaches to estimate the potential of C-sequestration of agricultural land. To quote some examples for Europe: First estimates of agricultural SCS potentials in Europe were given by Smith et al., (1998, 1996) based on several LTEs in Europe. Freibauer et al., (2004) conducted an update by combining other estimates by Batjes, (1996), Nabuurs et al., (1999), Smith et al., (2000 a,b), and Vleeshouwers and Verhagen, (2002) also taking into account the available surface of agricultural land and its suitability for implementation of those management options. Further estimates at the pan-European level were given by Lugato et al., (2014) using the CENTURY agroecosystem model for a set of management practices under different scenarios. Despite all these efforts, there is still an incomplete understanding of how SCS is influenced by climate, by land use and –management and by pedo-climatic factors (Chenu et al., 2019; Smith et al., 2020; Stockmann et al., 2013).

In contrast to mineral soils, effective mitigation measures for organic soils while maintaining intensive agricultural production are much less studied. Organic soils in peatlands are one of the most important terrestrial carbon sinks, storing as much carbon as all terrestrial biomass worldwide. When drained for



agriculture, they are a substantial source of greenhouse gases (GHG) due to the degradation of organic matter that is exposed to oxygen in the air. In the last decades, peatlands have been suffering degradation mainly through drainage for agriculture, forestry and peat extraction (Joosten and Clarke, 2002). Rewetting and restoration of cropped organic soils is one important option to mitigate GHG emissions, with great potentials. Globally ~ 1.91 (0.31–3.38) Gt CO₂-eq. yr⁻¹ could be saved by peatland restoration (Leifeld and Menichetti, 2018). An alternative way of using converted wet and rewetted peatlands for agriculture is paludiculture, which however comes at the expense of losing land for intensive food production (Leifeld and Menichetti, 2018).

In summary, it can be concluded that there are still essential knowledge gaps comprising the effectiveness and feasibility of mitigation options for both mineral and organic soils. Moreover, country-specific knowledge and data sets are hitherto insufficiently exploited and used for implementing mitigation projects in practice. With this stocktake we aim to provide a better overview on the available knowledge of achievable SCS potentials in mineral soils in agricultural land, including pasture/grassland across Europe, under different farming systems, soil types and pedo-climatic conditions as well as on GHG mitigation measures for managed organic soils. We present an overview of estimates on carbon sequestration (mineral soils) and greenhouse gas mitigation potentials (organic soils) and measures for agricultural soils. An overview of the share of land for which such information is available is given and finally we present the most urgent knowledge gaps regarding methodological approaches, soil or management types and geographical regions.

3. Methodology and data source

The stocktaking of achievable soil carbon sequestration is based on three pillars:

- Project organization, providing an overview of the published potentials including methods used, data coverage etc., and providing an overview of personnel and institutions involved.
- Project organization. AGS and CRAW identified the main experts to be approached by the various consortium members, with the help of information delivered by the project coordinator.
- To achieve the main scientific goal a questionnaire was developed by the subtask leaders and provided to experts of each MS who helped in gathering the information available on the internet or provided non-public reports and information to the subtask leaders from AGS and CRA-W (see list of MS experts and contacts in Annex D).



The questionnaire was divided in two parts: a) the request for providing information on the achievable SCS in mineral soils and b) the request for providing information on the potential for reducing GHG emissions from managed organic soils. (See Annex E for complete questionnaire) to which the results are assigned accordingly. Together with the literature and the questionnaire, the partners delivered a summary in text format with a maximum of three pages excluding references, which included the information on state-of-the-art knowledge on the achievable SCS in mineral soils in agricultural land, including pasture/grassland across Europe, under different farming systems, soil types and pedo-climatic conditions as well as on GHGs mitigation measures for managed organic soils. The contributions were gathered between June and the end of August 2020. Each contribution (questionnaire, summary and literature) was reviewed and analysed by the task leaders. Documents submitted by the MS that did not fulfil our criteria (e.g. studies not from the country, C-sequestration potentials of other land uses like forest) were excluded. Although implementation of climate change mitigation measures and agricultural practices is often a result of socio-economic decisions, SCS varies depending on different soil types and climate.




Figure 1 The Environmental Stratification of Europe in 84 strata. Where the size of the stratum permits, the individual strata are labelled within the main Environmental Zones. The stratification extends from 11° W to 32° E and from 34° N to 72° N. (Metzger et al., 2005).



The data was therefore primarily organized in respect to the boundaries of the Environmental Zones (EZ) as defined by Metzger et al., (2005) (default EJP SOIL) (Fig. 1). Additionally, results were also presented individually per MS where relevant. However, it is important to notice that implementation of agricultural mitigation measures is often guided by political, socio-economic and organizational considerations rather than by eco-physiological framework conditions. Consequently, the distribution of mitigation projects does not necessarily reflect the geographical variability of biophysical SCS potentials.

4. Results

4.1 Data and data coverage

The results presented here have to be taken with caution, as quality of the individual contributions was highly variable.

All participating MS (n= 25) returned their country specific information. These covered all EZ. For mineral soils, 153 contributions were considered for the stocktake. Out of them 80 % are published scientific literature and 20 % are grey literature (reports, presentations) (Fig. 2). Regarding the organic soils only 21 studies were considered with a share of 67 % published scientific literature and 33 % grey literature. All the literature dates from the years 2000-2021. More than 85 % of the articles were written in the period 2011-2021, reflecting the increase of interest in SCS over the last decade (Fig. 2). The lists of references for mineral and organic soils, respectively, grouped by MS are included in the Appendix A and B. In total 114 ongoing and planned research projects all over Europe were mentioned as being relevant for SCS and GHG-mitigation in agricultural soils (List in Appendix C).

The following results are presented separately for the mineral soils and organic soils, organized according to the questions posed to the MS's.

4.2 Mineral soils

SCS in mineral soils are covered by 153 contributions. The data density per EZ (studies per 10'000km²) was visualized on a map, counting the studies per EZ (Fig. 3). The area of the EZ was normalized by the agricultural area including arable land, permanent crops, grassland and heterogeneous agricultural areas. The agricultural area was extracted from raster data of the CORINE land cover (CLC) inventory of Europe 2018 using ArcGIS and overlaid with the EZ by Metzger et al. (2005). Looking at the distribution per EZ the, Alpin North (ALN), Nemoral (NEM) and Boreal (BOR) zones are best covered by studies while Mediterranean Mountains (MDM), Atlantic North (ATN), Lusitania (LUS), and pannonian



(PAN) (with decreasing density) are relatively poorly studied. The Atlantic Central zone (ATC) and the Continental zone CON, covering the greatest share of agricultural area are moderately to well studied.




Figure 2 Type of literature considering SCS and distribution from 1990-2021.

In total 38 different measures were identified and studied to evaluate SCS on mineral soils. To allow a quantitative evaluation of the results, the measures that were reported to have an effect on SCS have been summarized into five main groups: 1) land use (excl. forest), 2) soil protective measures, 3) tillage, 4) fertilizers and soil amendments, and 5) other. Under land use, we include all measures that imply a long-term change in the coverage of the land, which cannot be quickly reversed. Soil protective measures include all types of farming practices targeting or favouring soil protection such as permanent soil cover, crop rotation, crop residue management, other include agroforestry, organic farming etc. (see Table 1 for complete information). No-till practices, which generally fall into the soil protective category, were counted separately because of their great number. “Other” contain all measures, which do not fit into the defined groups.




Figure 3 Data density (Studies per 10'000km²) for each environmental zone (EZ) across Europe, normalized by its agricultural area. Value unit: Alpine North (ALN), Boral (BOR), Nemoral (NEM), Atlantic North (ATN), Alpine South (ALS), Continental (CON), Atlantic Central (ATC), Panonian (PAN), Lusitanian (LUS) Anatolian (ANA), Mediterranean mountains (MDM), Mediterranean North (MDN), Mediterranean South (MDS). EZ according to Metzger et al., 2005.



Table 1 Agricultural measures to improve SCS in mineral soils mentioned in the stocktake.

Measures	Total number of studies
1 Land use:	41
Permanent grassland	12
Conversion to grassland	10
Sown biodiverse permanent grasslands	2
Perennials	5
Agroforestry	9
Shrub encroachment	3
2 Soil protective measures:	82
Mulching	1
Ley farming	9
Cover crops	16
Catch crop	6
Improved crop rotation	16
Specific crops	1
Crop residue management	23
Grass residue management	1
Reduction of grassland renovation	5
Erosion control measures	1
3 Tillage:	81
No-till	35
Conservation tillage (minimum tillage, reduced tillage)	31
Non inversion tillage	10
Dual layer tillage	2
Deep ploughing	2
4 Fertilizers and amendments:	68
Mineral fertilizers	16
Manure application	30
Amendments:	
Mycorrhiza-inoculated compost	1
Compost of organic waste	10
Liming	3
Sewage sludge	1
Biochar	4
Biochar-blended compost	1
5 Other measures:	16
Managed field margins	2



Improved grazing management	2
Organic farming	4
Above and below ground biomass maize	1
Hedges	1
Extensive grassland management	1
Restoration of organic soils	1
Crops with improved N efficiency	1
Energy crops	3

4.2.1 Question 1 (Q1): Are there any quantitative estimations of the achievable carbon sequestration under different soil management in your country (finished or ongoing studies)?

In total 43 contributions were submitted of which 23 are scientific published articles, and 20 are national reports. Thirteen countries assessed the achievable SCS in agricultural soils at the national scale (BE, CH, DK, ES, FR, IT, IR, NL, NO, PL, PT, SE, UK), and three countries presented data for specific regions (BE, DE, IT). The potentials have been reported either as total potentials Tg C yr⁻¹ or Tg CO₂ yr⁻¹ or as annual rates Mg C ha⁻¹ yr⁻¹ and are summarized in Table 2. In order to realistically indicate the achievable SCS it is first essential to determine where and on which area a particular measure can be implemented. Further, costs and practical applicability of the measures should be considered. Studies considering all these conditions are scarce. Only one country (France) includes the additional costs for farmers and shows that in general the implementation of practices that sequester carbon results in a cost for the farmers (Pellerin et al., 2019). All other countries only include broad estimates of the areas on which measures technically can be applied without consideration of other factors (i.e., socio-economic parameters). Climate change has been considered by Italy and Poland only. The study of Mondini et al., (2012) clearly shows the importance of including climate change scenarios, because the response to compost application in Italy was only 0.13 t C ha⁻¹yr⁻¹ when considering climate change, which was three times lower than values (0.42 ha⁻¹yr⁻¹) estimated by Smith et al. (2008) and Freibauer et al., (2004), where climate change was not accounted for. Contrasting methodologies were applied by the different countries (for example, sampling depths ranging from 10 to 100 cm were considered), making the comparison of SCS potentials from one country to another difficult. For this reason, the shares (%) of the potentials to the yearly total domestic emissions from the agricultural sector (EA) were calculated when possible (Figure 4 and Table 2). EA was taken from the national GHG inventory reports (NIR) of each country submitted in 2020 ([National Inventory Submissions 2020 | UNFCCC](#)).

The total potential of each country for climate change mitigation varies greatly, ranging from 0.1 % to 28.5 % of EA (Fig. 3). ***It is however important to note that estimations are reported for a certain soil***



depth, area and time period and cannot be compared directly to each other (see Table 2 for details and references).

France, Italy, Poland and Spain report relatively high potentials of **5.8, 2.1, 1.6 Tg C yr⁻¹** and **2.9 Tg C yr⁻¹**, respectively. For France the potential corresponds to 28.5 % of EA, which is to be achieved by a combination of measures including agroforestry (Table 2). This potential is however associated with very high annual costs in the order of 2.3 billion euros. The reported potential for **Italy** could offset 25% of the EA by applying compost of organic waste to all the arable land in the country. For **Spain**, the potential covers 27 % of their EA, which is to be attained by the implementation of no-till. For **Poland**, this corresponds to 18 % of the EA, which is to be achieved by combining reduced tillage, crops residues and manure. It should be noted that this potential is only achieved if the total crop area of Poland is considered, which might not be realistic. A combination of cover crops, agroforestry, rewetting of drained organic soils and afforestation of set aside arable land would reduce 25.5 % of the national EA of **Sweden**. **Norway** reports biochar as the measure with the greatest potential (**0.245 Tg C yr⁻¹**) for storing carbon in Norwegian soil in a long-term perspective, which would offset 20 % of their GHG from the agricultural sector. However, it has to be noted that the production of biochar would rely on 50 % of the available biomass residues at the national scale and would therefore compete with other uses for this feedstock. Outstanding potentials for land use and land use changes are given by **UK** with **2.39 Tg C yr⁻¹** and by **Switzerland 0.245 Tg C yr⁻¹** covering 21 % and 16 % of their respective EA. **Belgium** would achieve a potential of **0.26 Tg C yr⁻¹** on cropland, which could offset 9.6 % of the EA by a combination of measures including forest recover (Table 2). The sum of all the reported potentials would amount **18.8 Tg C yr⁻¹**, which would offset 16 % of the total **European EA** (Table 2, Fig. 4).






Figure 4 Share of potentials to the annual total emissions from the agricultural sector for each country and share of measures involved. Dashed squares indicated unknown proportion of measures. Selection of maximum potential per country—additional potentials of countries are included in Table 2.



Table 2 Achievable domestic soil carbon sequestration (SCS) and reduction potential of the total emissions from the agricultural sector (EA) reported by the MS. PSL = Published scientific Literature, GL= Grey Literature.

Country	Spatial Scale	Area (ha) of applied measure	Temp. Scale	Measure	SCS-Potential Tg C/yr.	Percentage of total EA (%)	Total annual EA (Tg C)	Methods:	Reference:
Belgium	Regional (Flanders)	-	15	Permanent grassland, compost application, green manuring and management of crop residues (cereals)	0.05	1.8	2.7	Scenario analysis comparing change in an agricultural measure to 'business as usual' calculating SCS potential over period 2016-2030.	D'Hose and Ruyschaert, 2017 (GL)
	Regional (Flanders)	-	12	Green manuring, crop residue management, temporary pastures, organic farming and compost application	0.06	2.2	2.7	Increase of effective Organic carbon in the "current" (2002) and baseline (1990) year were calculated. (sampling depth:24 cm)	Sleutel et al., 2007 (PSL)
	National	458807	20	Bio-energy crops, farmyard manure, woodland restoration, no-till, cover crops, improved management of farmed peat soils, organic farming	0.2589	9.6	2.7	Estimates are based on literature values and assumptions on the area the practice can be applied.	Dendoncker et al., 2004
Denmark	National	2310000	26	cover crops, crops residue management, conversion to grassland	0.136	4.5	3.0	Modelling with C-Tool (Soil depth: 100 cm)	Taghizadeh-Toosi and Olesen, 2016 (PSL)
France	National	28500000	30	Cover crops, reduced till, organic manures conversion to grassland, hedges, reduction of mowing, grass cover in vineyards, intra plot agroforestry	5.7 (30 cm) 8.15 (100 cm)	27.2 (39)	20.4	Modelling (Soil depth: 100 cm) with the STICS and Pasim models	(Pellerin et al., 2019 (GL, English)
Germany	Regional (Bavaria)	3315800	-	Agroforestry, conversion to grassland, cover crops, improved crop rotation, organic farming	0.3-0.4	1.6-2.02	17.3	Estimate based on literature values and assumptions on the area the practices can be applied.	Wiesmeier et al., 2020 (PSL)
Ireland	National		10	Winter cover crops 0.51 Mg ha ⁻¹ yr ⁻¹ , winter cover crops combined with min. tillage 0.74 Mg ha ⁻¹ yr ⁻¹	0.08 – 0.1	1.3 – 1.6	5.2	Flux measurements in combination with modelling and LTE's. (Soil depth: 15 cm)	(Lanigan et al., 2012) (GL)
	National	351000	9	Reduced tillage (0.18-1 Mg ha ⁻¹ yr ⁻¹), crops residue management (0.44-0.6 Mg ha ⁻¹ yr ⁻¹).	0.11	2.2	5.2	Field experiments and modelling Roth and cohort model (Soil depth: 60 cm)	van Groenigen et al., 2011 (PSL)



	National	450,000	10	Improved management of grassland	0.26	4.9	5.2	Field experiments and modelling Roth and cohort model (Soil depth: 60 cm)	(Lanigan et al., 2018) (GL)
Italy	National	16284166	100	Compost of organic waste	2.1	25.6	8.2	Modelling with RothC. 12 climate scenarios	Mondini et al., 2012 (PSL)
	Regional (Foggia, Apulia Region)	505400	20	Compost of organic waste	0.031	0.38	8.2	Modelling with RothC10N and management scenarios	Bleuler et al., 2017 (PSL)
	Regional (Manfredonia, Apulia Region)	35454	20	Crop residues incorporation and water management	0.002 – 0.02	0.02-0.25	8.2	Modelling with RothC10N and management scenarios	Di Bene et al., 2016 (PLS)
Netherlands	National	-	20	Cover crops, managing field margins, No-till, non-inversion till, green manure, crop residue management, slurry, reduction of grassland renovation	0.27	5.4	4.9	Modelling with MITERRA-NL	Lesschen et al., 2012 (GL Dutch)
	National	298253	30	Improved crop rotation	0.245	4.9	4.9	Modelling with NDICEA version 6.2.1	Koopmans et al., 2019 (GL Dutch)
	National	298253		Reduction of grassland renovation	0.031	0.6	4.9	"	«
Norway	National	84000	?	Biochar	0.245	20.1	1.2	Estimate based on literature. Cover crops on 60 % of crop area.	Rasse et al., 2019 (GL)
	National	22400	?	Cover crops	0.057	4.7	1.2	Based on meta-analysis results by (Lehmann and Joseph, 2015)	Bøe et al., 2019 (GL)
Poland	National	10400000	20	Reduced tillage, crops residues, manure	1.6	18.3	9	Modelling: DNDC model	Faber and Jarosz, 2018 ((PSL, Polish)
Portugal	National	-	10	Sown biodiverse permanent pastures rich in legumes (SBPPR) (1.76 Mg ha ⁻¹ yr ⁻¹). No Area indicated.	-		1.9	Modelling (soil depth 10 cm)	Teixeira et al., 2011 (PSL)
Spain	National	7650588		No-till	2.9	26.9	10.8	Modelling Century and RothC, using C emission-sequestration rates coefficients of the IPCC.	Moreno-García et al., 2020 (PSL)
Sweden	National	1792000	30	Perennials, grassland, cover crops or catch crops	0.324	16.5	1.9	SOC change factors based on Swedish long-term trials	Wikström, 2019 (GL, Swedish)



	National	610000	25	Cover crops, agroforestry, afforestation of set aside land, rewetting of organic soils	0.5	25.5	1.9	-	Karlsson et al., 2020 (GL, Swedish)
Switzerland	National	920000	20	Grasslands	0.258 (max)	15.8	1.6	Estimate based on literature values and assumptions on the area these practices can be applied	Beuttler et al., 2019 (GL, English)
	National	10000	20	Deep ploughing	0.021	1.3	1.6	"	Beuttler et al. 2019 (GL, English)
	National	8230	40	No-till	0.0027	0.2	1.6	Estimate based on literature values. No-till	Leifeld et al., 2003 (GL, English)
UK	National	-	30	Grassland remaining grassland and cropland converted to grassland	2.39	21	11.2	UK Greenhouse Gas Inventory, 1990 to 2018	Brown et al., 2020 (GL)
Europe					Sum of all countries: 19	16	118.2	EU Emissions from agricultural Sector: 426. 473 Mt of CO ₂ equivalent	EAA report 2020



4.2.2 Measures and land use Q1

Soil protective measures have the highest occurrence in the reports of MS, representing 35 % of all reported measures (Fig. 5). This group of measures includes, in decreasing order of importance: crop residue management (n=9), cover crops (n=9), and improved crop rotation (n=2). With 27%, land use (change) is the second most mentioned group of measure, followed by tillage with 18%.

The total achievable SCS is commonly estimated by combining measures from all groups. Estimates of the achievable C-sequestration using just one single measure were provided for no-till (ES, DE, CH), cover cops (SE), perennial crops (SE) and biochar (NO).




Figure 5 Share of measures mentioned for national estimates of SCS. Question 1 (Qs1).

4.2.3 Methods Q1:

The time periods considered in the studies ranged from 9 to 40 years and averaged 20 years. In total seven studies report time period of less than 20 years. By definition, SCS related to a change in farming practices lasts only for a certain period, until soil carbon stocks reach a new steady state. The IPCC therefore recommends a time horizon of at least 20 years for estimates of SCS potentials. Findings from multiple experiments at Rothamsted (Hertfordshire, UK), however, show how rates of SCS can vary through time over the entire duration of long-term experiments (Poulton et al., 2018a) and that soils may reach a new C equilibrium only around 100 years after a land use change (Smith, 2014). The increase in soil carbon storage is not linear and secondly its rate decreases over time. This means that potentials as annual rates (Tg C year⁻¹) will be larger if only calculated for a short period (e.g. 10 or 20



years) and will decrease if calculated for a longer period (e.g. 100 years). This should be taken into account when comparing the reported potentials.

The most common approaches used to estimate the potential of SCS for agricultural land are modelling (for detailed information see the review of models by Campbell and Paustian, (2015), calibration of SOC stock change factors from local long-term experiments, scenario analysis, or application of SCS rates from the literature.

Norway, Switzerland and Belgium applied SCS rates from the literature to their agricultural areas. Sweden used SOC change factors based on long-term trials and Flanders calculated the increase of effective organic carbon in the “current” (2002) and baseline (1990) year. A variety of modelling approaches were used by DK, DE, ES, FR, IT, NL, PL and UK. The models mentioned are:

- **The MITERRA-NL** is an adapted model for Dutch specific data based on the European MITERRA-EU Model (Lesschen et al., 2012). C-dynamics under conversion of land use is modelled based on a Tier 2 approach according to IPCC guidelines.
- The **C-TOOL Model** was developed to enable simulations of the medium- to long-term changes in SOC in the profile (0-100 cm) of temperate mineral soils under agricultural management and was improved with specific data for Denmark (Taghizadeh-Toosi et al., 2014). C-TOOL applies a less complex structure, compared with other models, which facilitates more straightforward calibration. It requires only a few inputs (i.e., average monthly air temperature, soil clay content, initial soil C: N ratio and C inputs to the soil from plants and other sources).
- A combination of the **CENTRURY** and **RothC** model was used to calculate SCS under regional land use in Spain (Moreno-García et al., 2020). C emission-sequestration of available SCS studies comprising Spanish regions as well as coefficients from the Intergovernmental Panel on Climate Change (IPCC) were included.
- **RothC model** and its modified version **RothC10N** were used to calculate SCS in Italy at different spatio-temporal scale (Farina et al., 2018). These models require few easily available inputs (e.g., average monthly air temperature, precipitation, evapotranspiration, soil clay content, initial SOC, and C inputs to the soil from plants or other sources). RothC was used to perform a spatially explicit simulation of the potential increase in C storage in compost-amended agricultural soils at the national scale under a changing climate, while RothC10N to simulate



SOC dynamics at regional level under semiarid conditions in Southern Italy (i.e., Foggia Province and Manfredonia District).

- **The EPIC model** was used to estimate the CO₂-mitigation potential in the federal state of Baden-Württemberg (SW-Germany) comparing conventional and zero tillage and taking into account C losses through soil erosion (Gaiser et al., 2008). The adjusted net CO₂-mitigation rates of zero tillage, subtracting the reduced carbon losses through soil erosion, were estimated. In Italy, the EPIC model was used to estimate soil carbon dynamics at field level in a representative rainfed cereal system as influenced by climate change and contrasting tillage (Iocola et al., 2017).
- **The DNDC model** Denitrification - Decomposition (DNDC) model (version 9.2; <http://www.dndc.sr.unhu.edu>) is used to estimate changes of SOC in Poland (Faber and Jarosz, 2018). DNDC consists of six sub models for simulating the soil climate, plant growth, decomposition, nitrification, denitrification and fermentation. To simulate SOC changes in agricultural land, DNDC requires a number of input parameters including daily meteorological data (air temperature and precipitation), soil properties (bulk density and initial SOC context), and management practices (crop rotation, fertilization and manure application). DNDC has been tested against numerous field data sets of long-term change SOC and N₂O emissions at a regional and national scale worldwide.
- **The ICBM** (The Introductory Carbon Balance Model) was developed in Sweden. It is a dynamic model, which describes and predicts soil C dynamics in agricultural soils. It is a simple two-pool model (one young and one old C pool) (Andrén and Kätterer, 1997). It was developed with the aim as being a “minimum model” which includes only processes that are necessary and relatively well known in the way it can be understood and used by nonprofessionals. ICBM has been successfully applied to many LTE’s in different environments. An extended version ICBM/2 exists where two young pools, representing aboveground plant litter and roots are considered.



4.2.4 Question 2 (Q2): Are there any estimated C sequestration rates for different management practices derived from field studies in your country (finished or ongoing studies)?

In total 110 contributions, comprising local or regional field studies were submitted of which 100 are scientific published articles, two are PhD thesis and 10 are national reports.

4.2.5 Measures Q2

The share of measures mentioned for the field studies slightly differs from the measures used to estimate nationwide carbon sequestration. Most experiments include measures from the tillage group (31 %) followed by fertilizers and amendments group (28 %) and soil protective measures group (25 %) (Fig. 6). The most studied measures from the soil protective measures groups are crops residue management (n=15), improved crop rotation (n=14) and cover crops (=8). From the fertilizers group, manure application represents the largest share appearing in 30 studies. Soil protective measures, fertilizers or amendments are mostly combined. Land use cover 9% of the studies, pointing to a minor role when it comes to local and regional field experiments.




Figure 6 Share of measures mentioned for national estimates of SCS. Question 2 (Qs2).

4.2.6 Methods for quantifying changes in C stocks (Q2)

Three main points are essential when quantifying soil stock changes by comparing measurements taken at two or more points in time and by comparing different measures or land uses (Powlson et al., 2011, Smith et al., 2019):

- 1.) Top and subsoil should be sampled considering that measures can have opposite effects depending on the soil depth. For instance, the common positive effect of no-till on soil C content measured in the topsoil may not be apparent when measuring to 60 cm depth; also, crops with deep root systems require deep soil sampling (Haddaway et al., 2017; Smith et al., 2020).
- 2.) Any account for changes in soil bulk density should be considered when sampling whole soil profiles by using the equivalent soil mass or equivalent soil depth approach.
- 3.) The quantification of the grain size of fine earth (<2 mm) and quantification of the coarse mineral (>2 mm) fractions of the soil.

The IPCC recommends sampling to a minimum depth of 30 cm, which is not in accordance to the requirements listed above. The stocktake shows that 22 % of the studies consider less than 30 cm, 19 % sample precisely to 30 cm, 28 % sample deeper than 30 cm and 20 % do not mention the soil depth studied.

With regard to the tillage studies (reduced, minimum and no-till) which report soil depth, 16 % consider top soils down to less than 30 cm and 22 % precisely to 30 cm. For tillage experiments that account for the whole soil profile (eight studies), two studies include the equivalent soil mass approach.



Two out of nine studies report an increase in SCS using no-till (Farina et al., 2018, Lopez-Bellido et al., 2019) while seven state that there is no effect on C stocks when considering the whole soil profile (Anken et al., 2004; D’Hose et al., 2016; Dimassi et al., 2013; Hermle et al., 2008; Martínez et al., 2016; Meurer et al., 2018; Willekens et al., 2014). The two studies reporting an increase in SCS, however, refer to Mediterranean conditions and should be considered since the effect of no-till practices depend on pedo-climatic conditions (Chenu et al., 2019). In general, the rates provided by the MS for *Manure*, *Mineral fertilizers*, *No-till*, *Cover Crops* and *Residue management* (Fig.7) are comparable to values reported for meta-analysis of international and global studies (e.g. Freibauer et al., 2004; Poeplau and Don, 2015; P. Smith et al., 2000; Vleeshouwers and Verhagen, 2002).




Figure 7 Rates derived from provided studies: **Manure** (Baldi et al., 2018; Bleuler et al., 2017; Buysse et al., 2013; Fornara et al., 2016; Jokubauskaite et al., 2016; Kauer et al., 2019; Koopmans et al., 2019; Mondini et al., 2012; Singh and Lal, 2005; Taghizadeh-Toosi et al., 2014; Vanden Nest et al., 2015); **Mineral fertilizers** (Baldi et al., 2018; Borrelli et al., 2011; Dersch and Böhm, 2001; Fornara et al., 2016, 2013; Kauer et al., 2019; Keel et al., 2019; López-Bellido et al., 2020; Mazzoncini et al., 2016; Poeplau et al., 2018; Singh and Lal, 2005; Taghizadeh-Toosi et al., 2014); **No-till** (Anken et al., 2004; Barbera et al., 2012; Borges et al., 2018; Cillis et al., 2018; D’Hose et al., 2016; Dimassi et al., 2014; Freudenschuß, 2010; González-Sánchez et al., 2012; Hermle et al., 2008; Iocola et al., 2017; Keel et al., 2019; Leifeld et al., 2003; López-Bellido et al., 2020; Martínez et al., 2016; Mazzoncini et al., 2016; Meurer et al., 2018; Moreno-García et al., 2020; Powlson et al., 2012; Willekens et al., 2014); **Cover crops** (Autret et al., 2016; Bleuler et al., 2017; González-Sánchez et al., 2012; Mazzoncini et al., 2011; Poeplau and Don, 2015; Taghizadeh-Toosi et al., 2014; Wikström, 2019); **Residue management** (Buysse et al., 2013; Dersch and Böhm, 2001; Lesschen et al., 2012; Lugato et al., 2006; Poeplau et al., 2015; Poulton et al., 2018a; Singh and Lal, 2005; Taghizadeh-Toosi et al., 2014; Triberti et al., 2008).



The time periods mentioned for calculated carbon sequestration rates in the stocktake range from 2 to 100 years (mean 20 years). A total of 39 studies (40 %) consider time periods ≥ 20 years (IPCC default period) (Fig. 8).




Figure 8 Counts for time periods over which SCS was estimated.

4.3 Organic soils

Organic soils are one of the most important terrestrial carbon sinks, storing as much carbon as all terrestrial biomass in the world. When drained for agriculture, they are a substantial source of greenhouse gases (GHG). In the last decade, peatlands have been suffering degradation mainly through drainage for agriculture, forestry and peat extraction (Joosten and Clarke, 2002). Organic soils account for 10 % of the total agricultural area for participating MS, with 70 % located in northern Europe (Denmark, Finland, Norway, Sweden) (Tanneberger et al., 2017). According to estimates by Joosten & Clarke (2002), in total 57 % of this area is used for agriculture (DK 70%, FI 2%, SE 5%, NO 8%). Further large peatland areas, of which more than 70 % are used for agriculture, can be found in Germany, Netherlands and Poland. From 14 MS where organic soils comprise an essential area (>1000 km²) only seven MS presented data for GHG studies of organic soils (DE, IR, NL, SV, SE, CH and UK) and two (IT, SV) for alluvial organic rich soils in floodplains. National reports (incl. NIR) as well as scientific studies do mostly report emissions from organic soils but only few measures for mitigation of GHG emissions are presented. In total 21 contributions were delivered of which 14 are published scientific literature and 7 are grey literature.



4.3.1 Question 3 (Q3): Are there any quantitative estimates for reducing GHG emissions (CO₂, CH₄, N₂O) from managed organic soils under different soil management in your country (finished or ongoing studies)?

Only few publications give quantitative estimations of the potential for reducing GHG emissions (CO₂, CH₄, and N₂O) from managed organic soils. This is mainly due to a lack of scientifically based mitigation options that still allow intensive agricultural production.

Four countries report possible national mitigation potentials, all by rewetting of drained organic soils:

- **Ireland** reports a mitigation potential of **3.2 Tg CO_{2e} yr⁻¹** if 185,000 ha of drained histic soils were rewetted (50 % of the total drained histic soils in Ireland (Paul et al., 2018). This would compensate 16 % of the national GHG from the agricultural sector (EA) and 12 % also accounting for the EA inclusive LULUF sector. Emission savings were calculated by comparing emissions before and after rewetting. Generic (Tier 1) emission factors for organic soils (IPCC, 2014) were used to calculate emissions from drained and rewetted histic soils.
- **Switzerland** reports mitigation potentials of between **0.33 and 0.99 Tg CO_{2e} yr⁻¹** for an area of 17,000 ha (Leifeld et al., 2003). This would lead to a compensation of between 5.6 %-17 % of the national EA. Estimates are based on literature values for rewetting of drained soils and assumptions on the area these can be applied.
- In the **UK** 95,000 ha of peatland have been subject to some form of active restoration, of which around 70,000 ha involve some form of re-wetting. In total, these activities are estimated to have generated an emissions reduction since 1990 of **0.43 Tg CO_{2e} yr⁻¹** (Evans et al. 2017). For UK this had led to compensation of 1 % of their EA's.
- Indirect potentials are indicated by **Germany**, showing that high GHG emissions result from draining peatland for cultivation. On average, **7.5 Mg CO₂-C ha⁻¹ or 27.7 Mg CO₂ ha⁻¹ yr⁻¹** were measured on German grasslands which were established on drained organic soils (Tiemeyer et al., 2016). Thus, rewetting drained peatland provides a large GHG mitigation potential and might even cause small net C sequestration in the future.
- **The Netherlands** reports an annual loss from cultivated organics soils of **1.158 Mt** of C, respectively of **4.3 Mt of CO₂** emissions, which could be hindered by restoration of the organic soils. Further they present mitigation potentials, which are based on specific water management measures: water level fixation (200 kha), submerged drainage (80 kha), and



transition to paludiculture (20 kha), corresponding to **0,95 Mg CO_{2e} yr⁻¹**, **0,6 Mg CO_{2e} yr⁻¹** and **0,4 Mg CO_{2e} yr⁻¹**, respectively (Born et al., 2016). These results are based on the conversion of subsidence of the soil into CO₂ emissions. Direct GHG emissions are rarely measured. As a result, there are currently no experimental studies from the Netherlands that quantify the emission reduction potential on organic soils. Several national research programs and projects have however been initiated to study different effects of measures to prevent peat oxidation and improve predictions of GHG emissions (see list of projects in Annex C).

4.3.2. Question 4: Are there any estimated GHG mitigation potentials for managed organic soils derived from field studies in your country?

As already stated, there are still few studied mitigation options that still allow agricultural production. The **Netherlands** reports measures that include water level fixation by keeping the groundwater level constantly below 30-60 cm, drainage restrictions and alternate wetting and drying to be promising options to mitigate GHG. For instance keeping a constant groundwater level of 30-60 cm below ground can lead to a GHG reduction of 71.1 % or 17.8 Mg CO₂-eq/ha/yr (van den Akker et al., 2018). By comparing two grasslands in **Ireland** with contrasting drainage, soil nutrient status and management practices, Renou-Wilson et al. (2014) assessed the role of these parameters in respect to biomass productivity, GHG and waterborne C fluxes which lead to a net ecosystem C balance (NECB). The 2-year mean NECB of the deep- and shallow-drained nutrient-poor sites were considerably smaller than the nutrient-rich site (40 and 16 % respectively). Nutrient rich sites produce much higher GHG emissions and are hotspots of fluvial C losses. These results are relevant for the development of strategies to reduce GHG and the choice of grassland types in **Ireland**. Another study by Wilson et al. (2016) report that an estimated amount of **75 Mg CO₂-C a⁻¹** was mitigated by the restoration of the Bellacorick, bog complex (9519.75 ha) in east **Ireland**, this result in an estimated value of €1506 ha⁻¹ in avoided losses. Paul et al., 2018 reports that a considerable amount of **0,4 Tg CO_{2e} yr⁻¹** could be saved, if in nutrient rich grasslands drainage spacing was decreased to control the average water table at -25 cm or higher.

Ongoing **Swedish** field experiments are studying GHG mitigation measures such as sand addition, paddy and sphagnum farming. The first preliminary results for sand addition show that the CO₂ emissions is highest from the plots without sand addition (3.4 μmol*m⁻²s⁻¹) and lowest from the plots where 5 cm sand was added (1.4 μmol*m⁻²s⁻¹). The emission from the 2.5 cm treatment was 1.8 μmol*m⁻²s⁻¹. The sand was applied in the autumn of 2015 and mixed in the top 10 cm of the soil.



Penetration resistance, yield and CO₂ emissions will be compared during three years. Results will be available soon (see list of ongoing projects in Annex C).

In **Italy**, organic soils are relatively scarce (870 km²), corresponding to 0.29 % of the total area. The area of organic soils cultivated annually (histosols) is estimated to be 232.47 km², corresponding to 0.18 % of the utilized agricultural area (ISPRA, 2020). The N₂O emissions from these cultivated soils are negligible and they are estimated to be 290 tons in 2018, representing 0.82 % of N₂O emissions for the agricultural sector. Nevertheless, Italy is the largest rice-producing region in Europe with 2170 km², corresponding to 1.68 % of the utilized agricultural area in Italy (ISPRA, 2020), where rice cultivation is most widespread in the Po Valley (Northern Italy). Methane (CH₄) emissions from rice cultivation have been identified as a key source in Italy. In 2018, CH₄ emissions from rice cultivation were 62,122 Mg, corresponding to 1.55 Tg CO₂eq, and representing 8.1% of CH₄ emissions from the agricultural sector (equal to 5.1 % of total CO₂eq). In Northern Italy, rice seeds are generally mechanically broadcasted in flooded fields. However, in the last 15 years, the submersion of the rice paddy field has been replaced by rice direct sowed into dry soil in rows, frequently irrigated. In 2018, direct sowing into dry soils covered 50 % of the total area cultivated with rice; its emission factor (EF) was about 25 % lower compared to the EF of the submersion sowing (24.74 vs. 32.69 g CH₄ m⁻² yr⁻¹). In this context, some interesting studies involving GHG flux measurements comprising rice cultivation in organic-rich, but mineral alluvial floodplain soils have been carried out (Bertora et al., 2018; Lagomarsino et al., 2016; Meijide et al., 2017). Bertora et al. (2018) compare water management techniques such as seeding during flooding- or dry periods combined with tillage and crop residue management and show that the adoption of dry seeding resulted in the lowest cumulative CH₄ emissions with a reduction of 69 % compared to the water-seeding analogue. Meijide et al. (2017) investigate the effectiveness of managing the water table level of rice paddy fields and observe that an adequate management of the water table level has the potential to decrease the Global Warming Potential and reduces CH₄ fluxes. Similarly, Peyron et al. (2016) evaluate the effectiveness of alternative water management techniques (i.e., dry seeding with delayed flooding and intermittent irrigation) on the mitigation of CH₄ emissions from paddy soils and show that the adoption of alternative water management reduces annual CH₄ emissions by 60 % compared to the conventional water seeding and continuous flooding practices, while adoption of intermittent irrigation regimes can totally prevent CH₄ emissions. Lagomarsino et al. (2016) show how alternating wetting and drying reduces CH₄ but can significantly increase N₂O fluxes depending on the transition time from flooding to anaerobic conditions. These studies used chambers to measure GHG fluxes.



4.4 Knowledge gaps reported by MS

All member states (MS) received a targeted questionnaire aimed at compiling knowledge gaps regarding the achievable SCS in mineral soils and the potential for reducing greenhouse gas (GHG) emissions from managed organic soils. This report illustrates how the obtained information was processed in detail and presents the conclusions after synthesising all information received.

4.3.1 Knowledge gaps in C-Sequestration in mineral soils

In total, fifteen MS submitted 77 knowledge gaps regarding C sequestration in mineral soils in Europe. They were first structured by the formulated knowledge gap, whether it was published or not, and which MS stated the knowledge gap. Moreover, similarities between knowledge gaps were highlighted with the help of colour coding in order to identify connections throughout Europe. Based on this information, related knowledge gaps were grouped and formulated:

1. Measures for SCS
2. Assessment of SCS
3. Interaction of soil management approaches
4. Chemical, physical, biological and climatic factors
5. Economic, socio-economic and social factors

Briefly, most knowledge gaps stated by the MS concerned the impact of soil management on C sequestration potential of agricultural soils and the assessment of soil C pools. The number of knowledge gaps per group and the MS stating the gaps are presented in Table 3. The knowledge gaps for mineral soils are summarized for the European regions in Table 4. In the following, the findings, which were obtained by grouping the knowledge gaps, are presented in detail. Published studies are indicated with the respective reference.

Table 3 Counts of knowledge gap mentions per group and MS

Knowledge gap	Counts per group	Member states
Measures for SCS	27	BE, FR, IT, NL, NO, ES, SE, TR
Assessment of SCS	22	AT, BE, CH, FR, IE, IT, LT, NO, PT, ES, SE, TR, UK
Interaction of soil management approaches	10	AT, BE, CH, FR, IE, IT, NL, SE, UK
Chemical, physical, biological and climate factors	12	IE, FI, FR, ES, SE, UK, CH
Economic, socio-economic and social factors	6	BE, FR, IT, CH
total	77	



Table 4 Counts of knowledge gaps per European region

European Region	Member states per region	Counts per region
Northern Europe	NO, SE, FI, DK	19
Western Europe	IE, UK, NL, BE, FR	23
Central Europe	EE, LV, LT, PL, DE, CZ, AT, CH, SK, HU, SI	12
Southern Europe	PT, ES, IT, TR	23

Measures for SCS

The group “Measures for SCS” summarizes knowledge gaps about agricultural management options aiming at increasing soil carbon sequestration. This group of knowledge gaps is therefore closely connected with the EJP SOIL challenge “**Maintain/increase SOC**” (Fig. 9). Since participating MS defined different knowledge gaps related to this group, a detailed description for each country is presented.

Belgium (Flanders) and France report that the effect of grassland renewal/renovation on C stocks (reseeding) is understudied. Moreover, the effect of intensity of grassland management (number of cuts, grazing intensity) and species composition on C sequestration potential needs to be investigated. Detailed information on C input from above- and below-ground biomass residues and the contribution of roots of different crop plants and different cropping systems is largely missing as well. Limited understanding of carbon inputs by rhizodeposition was also noted as a knowledge gap by **Switzerland**. According to **Belgium (Wallonia) and France**, the potential of deep-rooting plants and agroforestry/agroecology practices in temperate context for SCS deserve closer attention. Subsoil carbon storage and persistence of sequestered SOC are other topics that are currently poorly addressed. Research on new processes to clean waste organic effluents (like sewage sludge or biosolids) and make them safe for recycling in agriculture is also critical to decrease the waste of exogenous organic matter.

Considering **Norway**, so far, no study on the use of different cover crops in relation to C sequestration has been published that is considered to cover this topic sufficiently. Especially data on the potential use of deep root cultivars is not available (and is generally limited internationally), even though they are considered to have a significant potential for C sequestration. Regarding conservation agriculture, there is a general knowledge gap on how much C sequestration potential can be obtained from this practice in Norwegian (C-rich) soils. **Norway** further highlights that the use of organic resources to increase SOC in agricultural soils is a major knowledge gap, especially regarding the net effect of using organic resources. The ability of biochar to influence the stabilization of plant-derived C and other C inputs needs to be looked into for Norwegian conditions (Rasse et al., 2019). Finally, optimized management of permanent grasslands is also an understudied topic in **Norway**.



Sweden also points out the lack of information on the net effect of biochar on GHG emissions and the net effect of irrigation on SOC stocks. The unknown fate of biochar in soil was also identified as a knowledge gap by **Switzerland and Belgium (Wallonia)**. Both recognise that biochar has a strong potential for SCS but also point out that the design of biochar application to soil has to primarily take into account its effect on soil fertility. As biochar lasts in soil for centuries to millennia, a precautionary approach is indispensable and in-depth study of the conditions of biochar application to soil for a successful use, as well as an evaluation of associated risks, is urgently needed. Moreover, the short-term effects of loosening and incorporation of straw slurry into the upper subsoil on soil physical properties and crop yield are understudied. Similar to Belgium and Norway, Spain considers the effects of management practices on C sequestration in grasslands as a knowledge gap.

In the **Italian** stocktake, the development of a landscape level assessment of the soil C sequestration rate in fruit orchards with poor SOC levels, starting with surveying the current management practices, is expressed as a knowledge gap. Italy also plans to extend its efforts towards understanding SCS potentials in different silvo-pastoral agroforestry systems and sheep-grazed pastures without the presence of trees. Further, few studies are examining the C stock in olive grove and vineyards in Italy. Assessing the dynamic changes of SOC stocks over time, using repeated sampling or chronosequence approaches, would allow to draw conclusions on the effect of soil management practices on SOC stocks in the long term. Optimization of cover crop management and the introduction of intercropping could contribute to enhance overall agroecosystem sustainability. Further studies including a larger number of sites and simulating SOC dynamics with modelling are necessary to achieve more general conclusions and to consider specific side-effects of contrasting tillage practices on SOC stock change.

In the **Netherlands**, measures whose effects are insufficiently substantiated to date are adjustments in crop rotations. Crop rotations including more cereal crops have been studied in 2018-2020, but including insufficient combinations so far. In addition, the application of green manure, grass-maize rotations, compost and organic manure in livestock farming and herb rich grasslands are understudied. The effect of additional measures introduced by the Common Agricultural Policy (CAP) also have not been quantified for their effect on SCS. This includes measures such as buffer strips, hedges and bird fields. **Spain** states that there is a need to assess SCS for managed grasslands (Aguilera et al., 2019).

Turkey concludes that research efforts have to be increased on pastures, marquis shrublands, wetlands, poplar and orchards, urban areas and agricultural areas. More information should be gained on the use of organic materials such as animal manure, biochar and compost, organic crops, crop fertilizers and green fertilization, which will increase the C stock and soil quality. **Switzerland** noted that it is yet unclear whether conversion to organic fertilizers gives a net benefit for SCS. In this regard,



it must be considered that the re-allocation of organic residues across units of land does not represent a SCS measure (Powlson et al. 2011, Schlesinger 2000).

Assessment of SCS

This group summarizes knowledge gaps about methodological approaches to determine soil carbon sequestration (SCS) and can be connected to the EJP SOIL knowledge gap “**Maintain/increase SOC**” (Fig. 9).

Austria states the need to develop a quantitative/semi-quantitative method to determine SCS potentials. There are already initial approaches, but further development of these first methodological approaches is still necessary in order to draw a more differentiated picture of future possibilities and to be able to assess the contribution of different measures to achieve the climate goals more realistically. For example, many Austrian publications do not measure but calculate bulk density of soil, which can lead to inaccurate results regarding C stocks or sequestration. This issue will be looked into within the ongoing CASAS project (Carbon Sequestration in Austrian Soils). Similarly, **Spain** also notes that bulk density is often not measured and that more long-term studies in a broader range of representative systems and information n about fixed soil mass accounting of SOC stock changes are needed (Aguilera et al., 2019, 2013).

Belgium (Wallonia) underlines that, comparably to the “4per1000” scenario projected in France, the gains in terms of SCS that can be expected from innovative agricultural practices should be simulated to help decision-makers taking efficient measures for climate change mitigation. In that goal, the mapping of the soil carbon saturation deficit is indispensable to identify agricultural areas with the highest SCS potentials and design regional or national management strategies accordingly. Such a map exists for Wallonia but several elements might contribute to decrease the uncertainty of the prediction and therefore increase the accuracy and reliability of the map.

In **Italy**, reliable reference values in terms of SOC stock for any land use, management, soil type and climate are stated as knowledge gaps. Moreover, SOC sequestration rates achievable under different pedoclimatic conditions by applying specific strategies of land management are understudied. SOC variation in the sub-soil (below 30 cm soil depth) and soil inorganic C pool size and variation in the top and in the sub-soil is insufficiently addressed. **Italy** expresses that more comparisons between soil respiration results on similar experiments as well as data on GHG emissions and mitigation strategies for a wide range of land use and cropping is needed. Moreover, Italy stated that it would be necessary to develop a landscape level assessment of the GHG mitigation potentials of climate smart practices in fruit orchards with poor SOC levels. To better identify the drivers of GHG mitigation potentials, it



will be important to pay attention to site-specific farming options, considering soil physicochemical properties and local water quality.

Ireland, Sweden and France (Pellerin et al., 2019) reported knowledge gaps on potential of subsoil for SCS as well. **Ireland** adds that the quantification of mechanisms for movement of OC into subsoils have to be better understood (Simo et al., 2019; Torres-Sallan et al., 2017). **Lithuania** expresses an insufficient integrated evaluation of SCS at the national scale and a general lack of research data on SOC stocks.

Norway has very limited information on SOC storage on a large part of the grasslands and rangelands, which make up a very large share of Norway's terrestrial ecosystems (Rasse et al., 2019)

The **Portuguese** national GHG inventory, which is the base for the estimation of the SCS potential, is not up to date. There is an urgent need for the extensive collection of data on soil C. This is clearly stated in the national inventory report where there is a recommendation for a detailed evaluation of SOC stocks to be carried out, together with the national inventory.

In **Switzerland**, there is a need to evaluate management practices that enhance SCS carried out at the regional or farm level, where site and management specific constraints can be considered (Leifeld et al., 2003).

In **Turkey**, there is a need for long-term measurements on SCS. They refer to the findings from multiple experiments at Rothamsted (Hertfordshire, UK) that show how rates of SCS can vary significantly through time during the entire duration of long-term experiments (Poulton et al. 2018).

Interaction of soil management approaches

This group focuses on knowledge gaps related to a combination of agricultural management options and their interactions. This section is in line with the EJP SOIL challenges “**Avoid N₂O/CH₄ emissions**” and “**Maintain/increase SOC**” (Fig. 9).

In **Austria**, GHG emissions from soils and SCS are rarely measured simultaneously, even though both are strongly affected by different management practices. Comparably, **Belgium (Wallonia)** considers that the complete life cycle analysis of agricultural systems representative of the regional context is needed to assess as well as possible their performance in terms of overall C balance and not only SOC storage or sequestration. A combined approach, estimating overall effects of farming practices on C dynamics is still missing. **Belgium (Flanders)** expresses that the relationship between permanent grassland age, ground water table level and C stocks is insufficiently investigated. Still, this knowledge could serve as a basis to pay farmers for keeping grasslands longer at the same place and for assessing C sequestration potentials.



Belgium (Flanders) states that an improved accuracy of regional N₂O simulations (using the DNDC model) is needed. Specifically, inclusion of more precise crop area surfaces, crop rotations, crop residues as well as improved simulations for temporary grassland, taking into account permanent grassland and spatial and temporal variations of N applications, are needed. There is a general need for high-quality N₂O emission datasets originating from field experiments including frequent N₂O measurements. New long-term N₂O emission datasets should account for differences in crop and grassland types, effects of land management, finer resolution for climate, geographical spread and temporal distributions. Further research focusing on soil-borne GHG mitigation strategies should include long-term field experiments and ensure a very frequent monitoring of N₂O in particular. The monitoring should take into account weather events and agricultural practices since these can strongly influence N₂O emissions. Finally, these datasets can facilitate the development of accurate models, which may allow better prediction of (regional) N₂O emissions. Therefore, in **Flanders**, GHG emission data on several crops and land management strategies are required.

More information on the (long-term) impact on N₂O emissions is required for:

- Slurry, farmyard manure, digestate and compost. To be precise, the application method, feedstock composition, timing of application, amount and frequency of application need to be studied more intensively.
- The effect of (reduced) tillage on GHG emissions, environmental circumstances during agricultural practices, the effect of specific crop rotations and catch crops and crop residue management.
- Nitrogen immobilization via addition of organic biological waste and crop residues: type of organic biological waste, application method and dose, and environmental circumstances.
- Emissions from temporary and permanent grassland and grazing vs mowing.
- Impact of land use change (including conversion of grassland to arable).
- Impact of temperature and moisture conditions in order to simulate seasonal and environmental conditions (including winter period) and effects of climate change.
- Life cycle-like assessments of soil management strategies fostering soil C sequestration, including a potential increase in mineral N fertilization and GHG losses during treatment and storage of organic materials prior to soil application.

France states that there is a general need to assess the effects of combinations of agricultural management options and their interactions is necessary to improve knowledge on SOC sequestration. The **Italian** stock-take states that assessing the effects of combining diversified and sustainable management practices on soil properties, such as the integration of no-till, cover crops, organic



fertilization, return of crop residues and crop rotation is necessary to improve knowledge on SOC sequestration. Moreover, the spatial variability of SOC stocks needs to be assessed considering soil management, cropping systems and environmental zones.

In the **Netherlands**, there is still a lack of knowledge about realistic estimates of the area where different measures can be applied and regarding the effects of multiple measures applied at the same time (i.e. whether measures add on or exclude one another if applied at the same time). Further, the trade-off between soil C sequestration and nitrous oxide emissions is seen as insufficiently investigated in **Sweden**. In the **United Kingdom**, no published studies have specifically addressed the interaction of multiple management practices (e.g. tillage, nutrient applications, and liming applications) on the soil C sequestration potential.

Switzerland states that there is an urgent need to combine several SOC enhancing practices to sequester a significant amount of C in Swiss agricultural soils. In addition, application of novel measures as biochar, deep soiling and agroforestry should be tested (Keel et al., 2019).

Chemical, physical, biological and climate factors

This group summarizes knowledge gaps about chemical, physical and biological factors that influence the kinetics of SOC mineralization and therefore may affect the residence time of SOC. It can be connected to EJP SOIL challenges “**Optimal soil structure**”, “**Enhance soil biodiversity**” and “**Enhance soil nutrient retention/use efficiency**” (Fig. 9).

Finland identified that the effect of increased SOC content on the leaching of phosphorus and dissolved organic C (DOC) and further impacts on surface water quality is unknown. **Spain** underlines that organic debris > 2 mm are not taken into account in the measurement of SOC as only soil < 2 mm is analysed. Accordingly, SOC stocks can be over- or underestimated depending on whether the coarse fraction > 2 mm is mainly composed of stones or made of organic residues, respectively. **Sweden** states the effect of Al and Fe on the stabilization of SOC as a knowledge gap as well as the effect of microbial communities on C use efficiency (CUE) and SOC stabilization (Meurer et al., 2020). Moreover, the effect of pH on SOC stabilization is insufficiently studied under Swedish conditions. Switzerland also highlighted that management effects on microbial CUE are largely unknown. The **United Kingdom** expresses the need for knowledge on how (through which biogeochemical mechanisms) the interaction of different management practices (e.g. tillage, nutrient applications, liming applications) ultimately affect SOC dynamics.

Ireland, France (Pellerin et al., 2019) and **Sweden** expressed the need for studies about the effect of climate change on achievable soil C sequestration and the SOC net balance. Sweden further states a knowledge gap about the major controls on priming and its net effect on the SOC balance.



Economic, socio-economic and social factors

Within this group, knowledge gaps related to economic and social factors were compiled. No direct connection to one of the EJP SOIL challenges could be drawn.

Developing an accurate and cost-effective system for C accounting at the farm level (including remote sensing, modelling and regional calibration) is considered as a knowledge gap in in **Belgium (Flanders)**. Efficient decision-support tools and extra knowledge are needed to choose the optimal strategies for organic materials that can be returned to the soil based on a complete life cycle analysis, avoiding negative trade-off. The **French** stocktake raises the question, whether the development of bioeconomy is likely to reduce the achievable SCS because of competing use for available biomass. Further, it states that the effect of storing practices on global GHG emissions are uncertain. In **Italy**, there is no public perception of the importance of SCS. **Switzerland** reports that the implementation of SCS activities would have severe consequences for the agricultural sector and characteristic appearance of the landscape. The socio-economic feasibility of possible measures is therefore still an open question (Leifeld et al., 2003).

Soil management for carbon	Maintain/increase SOC
Assessment of soil carbon	Maintain/increase SOC
Chemical, physical, biological and climate factors	No connection to a challenge
Interaction of soil management approaches	Maintain/increase SOC
Economic, socio-economic and social factors	Optimal soil structure Enhance soil biodiversity Enhance soil nutrient retention/use efficiency

Figure 9 Knowledge gap groups for mineral soils and the corresponding EJP SOIL challenges.

4.3.2 Knowledge gaps in organic soils

Twelve out of 24 MS submitted in total 33 knowledge gaps regarding organic soils in Europe (Table 5), which were separated into four groups:

1. Management options for organic soils
2. GHG emissions from organic soils
3. Assessment of organic soils and SOC
4. Other specify problems of organic soils



Table 5 Number of knowledge gaps for organic soils and member states per group

Group	Nr. of knowledge gaps per group	Member states within this group
Management options for organic soils	12	IE, NL, SE, DE, TR
GHG emissions from organic soils	9	BE, LT, LV, SE, CH, IT
Assessment of organic soils and SOC	7	AT, IE, IT, SE
Other specific problems of organic soils	3	AT, ES
Total	31	

Moreover, knowledge gaps were clustered by European regions (Table 6).

Table 6 Knowledge gaps for organic soils per main European region

European Region	Member states per region	Nr. Of knowledge gaps per region
Northern Europe	NO SE, FI, DK	6
Western Europe	IE, UK, NL, BE, FR	7
Central Europe	EE, LV, LT, PL, DE, CZ, AT, CHE, SVK, HU, SL,	11
Southern Europe	ES, TR	7

Management options for organic soils

This group comprises knowledge gaps about agricultural measures on peat soils and can be connected to the EJP SOIL challenge “Maintain/increase SOC” (Fig. 10).

The stock-take of the **Netherlands** showed that there is a lack of studies focusing on the assessment of GHG reduction potentials according to existing mitigation measures. For example, the influence of groundwater levelling needs to be quantified. Similar is stated by **Ireland** reporting that the impact of water table manipulation on emissions should be assessed. Another potential measure that has not been evaluated so far is the effect of paludiculture compared to intensively managed grasslands. **Sweden** stated that there is an urgent need for new management options to mitigate GHG emissions and erosion on agricultural organic soils since only a small fraction of the area can be rewetted. In addition, it would be important to study the effect on GHG emission in simple ways to regulate the ground water table by plugging open ditches in paludiculture and to follow the long-term effects of rewetting projects on GHG emissions. Lastly, Sweden expresses that, as a climate change adaptation measure, conversion of organic soils to biomass production sites must be investigated as a feasible alternative in dry years.

Several MS also formulated knowledge gaps regarding the effects of agricultural management measures on GHG emissions. **Germany** established that a coordinated re-wetting project database is necessary to evaluate management options on rewetted soils (e.g. paludiculture) and their effect on



GHG emissions. The effects of rewetting on the large diversity of organic soils should be studied, in particular fens (nutrient rich) and degraded organic soils. Swedish experts found that there is a need to measure the long-term effects of sand addition (or incorporation of underlying mineral material) and liming on GHG emission rates and C capture efficiency. As proposed by **Turkey**, long-term organic and inorganic fertilizer effects on CO₂ fluxes should be investigated.

GHG emissions from organic soils

This group summarizes knowledge gaps about GHG emissions from peat soils. It is related to the EJP SOIL challenge “Avoid N₂O/CH₄ emissions” (Fig. 10).

Belgium (Flanders) states that there is a general need for high-quality N₂O emission datasets originating from field experiment including frequent N₂O measurements (Roelandt et al., 2007). New long-term N₂O emission datasets should account for difference in a.) Crop and grassland types, b.) Effects of land management and c.) Finer climate, spatial and temporal resolution (Roelandt et al., 2005). Finally, high quality GHG emission (especially for N₂O) data sets for several crops and land management strategies are required for Flanders. **Italy** suggests the creation of a free available library for GHG emission factors. **Lithuania** and **Latvia** sees, despite qualified calculations in its National Inventory Report (NIR), knowledge gaps on experimental GHG emission data on organic and mineral soils and an absence of emission factors at national scale. In **Sweden**, there is a great need for continuous measurements of N₂O emissions in field trials with different mitigation options to capture spring and autumn peaks and to increase the knowledge on a process level. They suggest that such measurements could be combined with microbial studies. More emission data for organic soils at different spatial and temporal scales to assess the huge variation in emission rates within fields, between fields, between soil types, and between years is needed. Remote sensing techniques might be a suitable option for increasing our knowledge in this regard. In **Switzerland**, emission factors of organic soils are seen as knowledge gaps as well.

Assessment of organic soils and SOC

This section gathers knowledge gaps on mapping peat soils and measuring GHG emissions from peat soils, which is in line with EJP SOIL challenges “Maintain/increase SOC” and “Avoid peat degradation” (Fig. 10).

Austria identified that the mapping and reporting of peatlands was lacking in their national inventory, with no data on depth and bulk density of peat soils. **Switzerland** expressed similar knowledge gaps, also with respect to the amount of C stored in these soils (Wüst-Galley et al., 2020). **Italy** reported a lack in the availability of updated land use maps and georeferenced data about land management



practices as well as analysing, accounting, and mapping GHG fluxes at a landscape scale. **Ireland** stated there is a need for assessment of the drainage status of organic soils.

Other specific problems of organic soils

Knowledge gaps that could not be assigned to one of the groups mentioned above were collected within the group “Other specific problems of organic soils” (Fig. 10).

Austria mentioned that the definition of peatlands is an issue. Peatlands are generally seen as natural biotopes whereas managed peat soils without vegetation are often overseen, which is why publications and their results divert substantially. Moreover, estimates of GHG emissions from peatlands in the Alpine foothills are generally extrapolated thanks to data from other countries (e.g. Germany). The suitability of calibration data must be verified for this context. **Spain** recognizes a huge gap between research on mineral soils and organic soils, mineral soils being much more studied. Additionally, cropping systems have been studied for GHG mitigation/quantification purposes much more frequently than livestock systems. Organic soils in the north of the country (e.g. grasslands) belong to the latter. More efforts are necessary in this field and type of systems/soils.

Generally, it can be concluded that very few studies deal with GHG emission factors of specific agricultural management methods on organic soils in Europe. Knowledge gaps on this topic are diverse, ranging from methodological problems such as mapping peatland and correctly estimating their SOC contents, to the GHG mitigation potentials of specific management measures. The general lack of research on organic soils is also reflected by the number of MS who did not submit studies and reports regarding reduction of GHG emission from organic soils despite the significant occurrence of peatlands on their territory.

Management options for organic soils	Maintain/increase SOC
GHG emissions from organic soils	Avoid N ₂ O/CH ₄ emissions
Assessment of organic soils and SOC	Maintain/increase SOC Avoid peat degradation
Other specific problems of organic soils	No connection to a challenge

Figure 10 Knowledge gap groups for organic soils and the corresponding EJP SOIL challenges



4.5 General knowledge gaps and hot topics

This part of the report provides thorough information about the knowledge gaps and hot topics, which were identified throughout the synthesizing process of the national data submitted by the MSs. These are summarized into different main topics.

4.5.1 GHG mitigation measure of organic soils:

Although farmed organic soils in most European countries represent a small part of the total agricultural area, these soils contribute significantly to national greenhouse gas budgets. However, very few mitigation options exist to mitigate GHG emissions from organic soils without compromising agricultural production. Most GHG mitigation practices reported by the MS involve the restoration of organic soils, which means abandonment of land from food production or even any agricultural use. Only one contribution (NL) reports possible mitigation potentials, which are based on specific water management measures (water level fixation). Nevertheless, there is an increasing awareness of the need of mitigation measures reflected by the several ongoing research projects on peatland management (see list of research projects).

4.5.2 General methodological challenges:

The stocktake has shown that the comparability of results is challenging due to different methodological approaches implemented. One possible explanation is that most studies and experiments were initially designed for other purposes (soil fertility, yields etc.) than measuring SCS. There is a large variation in sampling depth and periods over which SCS is estimated. Soil bulk density is often missing and therefore estimated, which however can lead to wrong numbers. To harmonize methodologies and improve the comparability of studies, the IPCC makes useful recommendations for the assessment of SCS. Nevertheless, a standard protocol is still missing.

Many studies do include crops residues, organic compost or manure, but the alternative fate of the amendments are rarely discussed. Even though such amendments can lead to an increase in soil carbon, it does not automatically imply an additional net transfer of C from the atmosphere into the soil (see definition of SCS section 1)(Chenu et al., 2019; Powlson et al., 2011; Schlesinger, 2000).

4.5.3 Socio-economic factors:

Socio-economic factors are key for a successful implementation of long-term measures (Amundson and Biardeau, 2018; MacLeod et al., 2010). There is an essential need of information about feasibility of implementation of agricultural measures with a high SCS potential, including the economic



efficiency and social acceptability on a country scale. The stocktake shows that socio-economic factors are rarely included in the studies. For instance, consideration of a realistic area where measures can be implemented, practical applicability and costs are generally not taken into account.

4.5.4 SCS under climate change

There is an urgent need to consider the impact of climate change on agriculture to correctly design achievable SCS scenarios. At present, there are very few simulations of SCS conducted under different climate scenarios. Achievable carbon sequestration, as well as greenhouse gas emissions from the agricultural sector, must be considered in the context of climate change, which are expected to significantly affect soil carbon dynamics, land use, production systems and farming practices in a near future. The study of Mondini et al., (2012) clearly shows the importance of including climate change scenarios, which showed that sequestration rates could be three times lower than expected because of climate change. At present, almost no studies do include climate change scenarios for their estimates. Climate change will unevenly affect European regions and hence soil carbon dynamics (Kovats et al., 2014; Meersmans et al., 2016). Spatial predictions of future soil organic carbon stocks under different climate scenarios could be an important starting point for further investigation in key regions with highest SOC losses.

4.5.5 Trade off C-Sequestration vs other GHG:

A widely discussed knowledge gap is the trade-offs between SCS and other GHG emissions (Lugato et al., 2018; Paustian et al., 2016; Reay et al., 2012). There is a general agreement that long-term experiments should include a constant monitoring of GHG emissions, in particular of N₂O since it represents the largest source and is very dynamic in space and time (Lugato et al. 2018, Reay et al., 2012). Lugato et al. (2018) and Guenet et al., (2021), for instance, show that mitigation potential of soil carbon management can be significantly overestimated by neglecting N₂O emissions. The stocktake confirms that for all MS GHG emissions from soils and soil C sequestration are rarely measured simultaneously, even though they are both strongly affected by different management practices. There is one recent study by Launay et al. (2021) which take into account several factors including GHG balance, biomass production and nitrogen- and water-related impacts in addition to soil carbon stock changes. They use a high-resolution modelling approach and show that current systems in France, even though they are storing some C in soils, on average are strong sources of GHG. Finally, it highlights some key management issues to improve GHG balances.



4.5.6 Measures and alternatives:

In recent years, **biochar** has been promoted as C negative emission technology including positive effects on soil fertility and crop production (Smith, 2016; Lehmann, 2006). Soil amendment with biochar is mentioned by several MS as a farming practice that could have large potential for SCS. Biochar also has positive interactions with the soil N cycle, as it does not require large amount of fertilizer N to sequester C in soils and contributes directly to reducing N₂O emissions (Guenet et al., 2021b). However, biochar is a technology under development, and questions of productions costs and competition for feedstock with other use such as bioenergy need to be addressed. There is also a need to optimize the technology for simultaneous C sequestration gains and increases in crop productivity, which requires further research and development. Studies indicate that biochar produced under the right conditions for agricultural purposes is safe, and the existing strict certification processes need to be applied to guarantee that these conditions are met. In this context, an in-depth risk evaluation of existing studies is urgently needed to precisely define which feed stocks and production methods are appropriate.

Deep ploughing, a method used to improve soil structure or overcome hardpans of podzols, has been shown to increase SOC stocks by very significant amounts (Alcántara et al., 2016; Schneider and Don, 2019). Preliminary estimates made for Switzerland show that by applying deep ploughing on an area of 5000 ha per year, could offset the emissions of 15.4 million t CO₂ over 20 years. This would compensate 12 % of the annual emission from the agricultural sector. Nevertheless, it is important to note that deep ploughing is a major irreversible soil intervention influencing several soil ecosystem services (either positively or negatively), which has to be taken into account when optimising soil fertility for crop production (Schiedung et al.2019).

Deep soil carbon: Concerning deep soil horizons, it has been shown that the residence time of subsoil C can be several times higher than that of the topsoil. However, the potential for additional C sequestration in deep soil layers remains to be assessed (Rumpel and Kögel-Kanbner, 2010; Chenu et al., 2019) as well as the mechanisms controlling movement of OC into subsoils and its stabilization (Balesdent et al., 2018; Kögel-Knabner and Amelung, 2021; Simo et al., 2019; Torres-Sallan et al., 2017).

4.5.7 Soil (bio-) chemical properties

Most studies focussing on soil processes and farming practices that increase soil C storage report measurements of SOC content together with other soil fertility parameters such as total N content, cation exchange capacity physical soil properties (e.g. grain size, stone content). The mineralogy and



elemental composition of soils, as well as soil (micro-) biology are rarely considered. Therefore, the way these variables affect SOC stabilization is not clear yet (Lal, 2018). It has however been shown that carbon stocks are soil type specific (Kögel-Knabner and Amelung, 2021) and that interactions of geochemical factors and climate do influence soil organic carbon storage and turnover which should be considered for further soil carbon stock predictions (Doetterl et al., 2015).

4.5.8 Carbon use efficiency in differently managed systems

Carbon use efficiency (CUE), i.e., the ratio of carbon transformed to microbial biomass to carbon uptake by the decomposer organisms, is key to understanding soil carbon sequestration. Only subtle differences in CUE have large effects on soil carbon storage. CUE is related to many environmental and microbial parameters (e.g. temperature, pH, nutrient availability, decomposer community etc.) (Sinsabaugh et al., 2013) and might thus be manageable, thereby providing ways to more sustainable agriculture. However, how management actually influences CUE still needs to be explored.

6 Conclusions

All participating MS (n= 25) returned their country specific information. The stocktake shows that there has been a clear increase in interest for topics covering soil carbon sequestration for climate change mitigation in the last two decades. A total of 101 ongoing projects have been reported dealing with SCS and GHG mitigation measures. However, at present country based knowledge and engagement is still poor. Only half of the MS can provide information on achievable carbon sequestration at national and regional scales. Information provided is mostly based on rough estimates without consideration of technical and socio-economic feasibilities. Climate change is usually not included, although it is known that it will have an important impact on carbon stocks. The presented national SCS potentials do however point towards important contributions to mitigate climate change by covering considerable shares of national greenhouse gas emissions from the agricultural sector in the range of 1-28.5 %, underpinning the need for further and detailed investigations. The stocktake also shows that the availability of datasets concerning SCS is variable among Europe. While environmental zones in northern Europe and central Europe are relatively well studied, there is a lack of studies comprising parts of Southern, Southeastern and Western Europe. In contrast to mineral soils, effective mitigation measures for organic soils while maintaining agricultural production are much less studied. Very few mitigation options exist to mitigate GHG emissions without compromising agricultural production. Most GHG mitigation practices reported by the MS involve the restoration of organic soils, which



means abandonment of land from agricultural use. Finally, the comparability of results is challenging due to different methodological approaches implemented.

7 Limitations of the synthesis

The aim of this stocktake is to get an overview on the available knowledge of achievable carbon sequestration potentials in mineral soils in agricultural land, including pasture/grassland across Europe, under different farming systems, soil types and pedo-climatic conditions as well as on GHGs mitigation measures for managed organic soils. The results in this report have however to be taken with caution, as quality of the individual contributions was highly variable.

Specific limitation of the analysis arise from the following points:

- The level of detail of the information provided by the participating countries was not uniform. Some participants were quite extensive answering all the questions while other provided incomplete or no answers.
- The participating teams were likely to adopt different definitions to soil carbon sequestration. This is mainly because many studies providing information on soil carbon sequestration were not specifically designed to report carbon sequestration but soil fertility including data on soil carbon changes.
- The questionnaire could have been improved by a higher level of detail and restrictions for possible answers, which would lead to more homogenous answers more adequate for this analyse.

References

- Aguilera, E., Gaona, C.D., Palomo, C.R., Laureano, R.G., Rodríguez, M.S., Estévez, V.R., 2019. Producción ecológica mediterránea y cambio climático: estado del conocimiento (olivar). *Mercacei Mag.* 190–196.
- Aguilera, E., Lassaletta, L., Gattinger, A., Gimeno, B.S., 2013. Managing soil carbon for climate change mitigation and adaptation in Mediterranean cropping systems: A meta-analysis. *Agric. Ecosyst. Environ.* 168, 25–36. <https://doi.org/10.1016/j.agee.2013.02.003>
- Akker, J., Hoving, I., Hendriks, R., Knotters, M., 2018. . Onderwaterdrains zijn effectief (No. 2922). (Research report). Wageningen Environmental Research, Wageningen.
- Alcántara, V., Don, A., Well, R., Nieder, R., 2016. Deep ploughing increases agricultural soil organic matter stocks. *Glob. Change Biol.* 22, 2939–2956. <https://doi.org/10.1111/gcb.13289>
- Amundson, R., Biardeau, L., 2018. Opinion: Soil carbon sequestration is an elusive climate mitigation tool. *Proc. Natl. Acad. Sci.* 115, 11652–11656. <https://doi.org/10.1073/pnas.1815901115>
- Andrén, O., Kätterer, T., 1997. Icbm: The Introductory Carbon Balance Model for Exploration of Soil Carbon Balances. *Ecol. Appl.* 7, 1226–1236. [https://doi.org/10.1890/1051-0761\(1997\)007\[1226:ITICBM\]2.0.CO;2](https://doi.org/10.1890/1051-0761(1997)007[1226:ITICBM]2.0.CO;2)
- Anken, T., Weiskopf, P., Zihlmann, U., Forrer, H., Jansa, J., Perhacova, K., 2004. Long-term tillage system effects under moist cool conditions in Switzerland. *Soil Tillage Res., Soil Quality as an*



- Indicator of Sustainable Tillage Practices 78, 171–183.
<https://doi.org/10.1016/j.still.2004.02.005>
- Autret, B., Mary, B., Chenu, C., Balabane, M., Girardin, C., Bertrand, M., Grandeau, G., Beaudoin, N., 2016. Alternative arable cropping systems: A key to increase soil organic carbon storage? Results from a 16 year field experiment. *Agric. Ecosyst. Environ.* 232, 150–164.
<https://doi.org/10.1016/j.agee.2016.07.008>
- Baldi, E., Cavani, L., Margon, A., Quartieri, M., Sorrenti, G., Marzadori, C., Toselli, M., 2018. Effect of compost application on the dynamics of carbon in a nectarine orchard ecosystem. *Sci. Total Environ.* 637–638, 918–925. <https://doi.org/10.1016/j.scitotenv.2018.05.093>
- Balesdent, J., Basile-Doelsch, I., Chadoeuf, J., Cornu, S., Derrien, D., Fekiacova, Z., Hatté, C., 2018. Atmosphere–soil carbon transfer as a function of soil depth. *Nature* 559, 599–602.
<https://doi.org/10.1038/s41586-018-0328-3>
- Barbera, V., Poma, I., Gristina, L., Novara, A., Egli, M., 2012. Long-term cropping systems and tillage management effects on soil organic carbon stock and steady state level of C sequestration rates in a semiarid environment. *Land Degrad. Dev.* 23, 82–91.
<https://doi.org/10.1002/ldr.1055>
- Batjes, N.H., 1996. Total carbon and nitrogen in the soils of the world. *Eur. J. Soil Sci.* 47, 151–163.
<https://doi.org/10.1111/j.1365-2389.1996.tb01386.x>
- Bellamy, P.H., Loveland, P.J., Bradley, R.I., Lark, R.M., Kirk, G.J.D., 2005. Carbon losses from all soils across England and Wales 1978–2003. *Nature* 437, 245–248.
<https://doi.org/10.1038/nature04038>
- Bertora, C., Cucu, M.A., Lerda, C., Peyron, M., Bardi, L., Gorra, R., Sacco, D., Celi, L., Said-Pullicino, D., 2018. Dissolved organic carbon cycling, methane emissions and related microbial populations in temperate rice paddies with contrasting straw and water management. *Agric. Ecosyst. Environ.* 265, 292–306.
- Beuttler, C., Keel, S.G., Leifeld, J., Schmid, M., Berta, N., Gutknecht, V., Wohlgemuth, N., Brodmann, U., Stadler, Z., Tinibaev, D., Wlodarczak, D., Honegger, M., Stettler, C., 2019. The Role of Atmospheric Carbon Dioxide Removal in Swiss Climate Policy – Fundamentals and Recommended Actions., Report by Risk Dialogue Foundation. Commissioned by the Federal Office for the Environment, Bern., Bern.
- Bleuler, M., Farina, R., Francaviglia, R., di Bene, C., Napoli, R., Marchetti, A., 2017. Modelling the impacts of different carbon sources on the soil organic carbon stock and CO₂ emissions in the Foggia province (Southern Italy). *Agric. Syst.* 157, 258–268.
<https://doi.org/10.1016/j.agsy.2017.07.017>
- Bøe, F., Bechmann, M., Øgaard, A.F., Sturite, I., Brandsæter, L.O., 2019. Fangvekstenes økosystemtjenester – Kunnskapsstatus om effekten av fangvekster (Nibio No. 5). Norway.
- Borges, O., Raimundo, F., Coutinho, J., Gonçalves, B., Oliveira, I., Martins, A., Madeira, M., 2018. Carbon fractions as indicators of organic matter dynamics in chestnut orchards under different soil management practices. *Agrofor. Syst.* 92, 301–310.
- Born, G.J. van den, Kragt, F., Hensens, D., Rijken, B., Bommel, B. van, Sluis, S. van der, Polman, N., Bos, E.J., Kuhlman, T., Kwakernaak, C., Akker, J. van den, Diogo, V., Koomen, E., Lange, G. de, Bakel, J. van, Brinke, W.B.M. ten, 2016. Dalende bodems, stijgende kosten: mogelijke maatregelen tegen veenbodemdaling in het landelijk en stedelijk gebied: beleidsstudie. Planbureau voor de Leefomgeving.
- Borrelli, L., Colecchia, S., Troccoli, A., Caradonna, S., Papini, R., Ventrella, D., Dell’Abate, M.T., Galeffi, C., Tomasoni, C., Farina, R., 2011. Effectiveness of the GAEC standard of cross compliance Crop rotations in maintaining organic matter levels in soils. *Ital. J. Agron.* 6, 8.
<https://doi.org/10.4081/ija.2011.6.s1.e8>
- Brown, P., Cardenas, L., Choudrie, S., Jones, L., Karagianni, E., MacCarthy, J., Passant, N., Richmond, B., Smith, H., 2020. UK Greenhouse Gas Inventory, 1990 to 2018: Annual Report for



- submission under the Framework Convention on Climate Change - Defra, UK. Department for Environment, Food and Rural Affairs (Defra), Nobel House, 17 Smith Square, London SW1P 3JR helpline@defra.gsi.gov.uk.
- Buyse, P., Roisin, C., Aubinet, M., 2013. Fifty years of contrasted residue management of an agricultural crop: Impacts on the soil carbon budget and on soil heterotrophic respiration. *Agric. Ecosyst. Environ.* 167, 52–59. <https://doi.org/10.1016/j.agee.2013.01.006>
- Campbell, E.E., Paustian, K., 2015. Current developments in soil organic matter modeling and the expansion of model applications: a review. *Environ. Res. Lett.* 10, 123004. <https://doi.org/10.1088/1748-9326/10/12/123004>
- Chenu, C., Angers, D.A., Barré, P., Derrien, D., Arrouays, D., Balesdent, J., 2019. Increasing organic stocks in agricultural soils: Knowledge gaps and potential innovations. *Soil Tillage Res., Soil Carbon and Climate Change: the 4 per Mille Initiative* 188, 41–52. <https://doi.org/10.1016/j.still.2018.04.011>
- Cillis, D., Maestrini, B., Pezzuolo, A., Marinello, F., Sartori, L., 2018. Modeling soil organic carbon and carbon dioxide emissions in different tillage systems supported by precision agriculture technologies under current climatic conditions. *Soil Tillage Res.* 183, 51–59.
- Dendoncker, N., Van Wesemael, B., Rounsevell, M.D.A., Roelandt, C., Lettens, S., 2004. Belgium's CO₂ mitigation potential under improved cropland management. *Agric. Ecosyst. Environ.* 103, 101–116. <https://doi.org/10.1016/j.agee.2003.10.010>
- Dersch, G., Böhm, K., 2001. Effects of agronomic practices on the soil carbon storage potential in arable farming in Austria. *Nutr. Cycl. Agroecosystems* 60, 49–55.
- D'Hose, T., Ruyschaert, G., 2017. Mogelijkheden voor koolstofopslag onder gras- en akkerland in Vlaanderen. (No. No. 231), ILVO Mededeling. Netherlands.
- D'Hose, T., Ruyschaert, G., Viaene, N., Debode, J., Vanden Nest, T., Van Vaerenbergh, J., Cornelis, W., Willekens, K., Vandecasteele, B., 2016. Farm compost amendment and non-inversion tillage improve soil quality without increasing the risk for N and P leaching. *Agric. Ecosyst. Environ.* 225, 126–139. <https://doi.org/10.1016/j.agee.2016.03.035>
- Di Bene, C., Marchetti, A., Francaviglia, R., Farina, R., 2016. Soil organic carbon dynamics in typical durum wheat-based crop rotation of Southern Italy. *Ital. J. Agron.* 11. <https://doi.org/10.4081/ija.2016.763>
- Dimassi, B., Cohan, J.-P., Labreuche, J., Mary, B., 2013. Changes in soil carbon and nitrogen following tillage conversion in a long-term experiment in Northern France. *Agric. Ecosyst. Environ.* 169, 12–20. <https://doi.org/10.1016/j.agee.2013.01.012>
- Dimassi, B., Mary, B., Wylleman, R., Labreuche, J., Couture, D., Piraux, F., Cohan, J.-P., 2014. Long-term effect of contrasted tillage and crop management on soil carbon dynamics during 41 years. *Agric. Ecosyst. Environ.* 188, 134–146. <https://doi.org/10.1016/j.agee.2014.02.014>
- Doetterl, S., Stevens, A., Six, J., Merckx, R., Van Oost, K., Casanova Pinto, M., Casanova-Katny, A., Muñoz, C., Boudin, M., Zagal Venegas, E., Boeckx, P., 2015. Soil carbon storage controlled by interactions between geochemistry and climate. *Nat. Geosci.* 8, 780–783. <https://doi.org/10.1038/ngeo2516>
- Faber, A., Jarosz, Z., 2018. Modelowanie emisji podtlenku azotu i amoniaku w skali regionalnej oraz w Polsce. *Zesz. Nauk. SGGW W Warszawie - Probl. Rol. Świat.* 18(33), 70–81. <https://doi.org/10.22630/PRS.2018.18.2.35>
- Farina, R., Testani, E., Campanelli, G., Leteo, F., Napoli, R., Canali, S., Tittarelli, F., 2018. Potential carbon sequestration in a Mediterranean organic vegetable cropping system. A model approach for evaluating the effects of compost and Agro-ecological Service Crops (ASCs). *Agric. Syst.* 162, 239–248. <https://doi.org/10.1016/j.agsy.2018.02.002>
- Fornara, D.A., Banin, L., Crawley, M.J., 2013. Multi-nutrient vs. nitrogen-only effects on carbon sequestration in grassland soils. *Glob. Change Biol.* 19, 3848–3857.



- Fornara, D.A., Wasson, E.-A., Christie, P., Watson, C.J., 2016. Long-term nutrient fertilization and the carbon balance of permanent grassland: any evidence for sustainable intensification? *Biogeosciences* 13, 4975.
- Freibauer, A., Rounsevell, M.D.A., Smith, P., Verhagen, J., 2004. Carbon sequestration in the agricultural soils of Europe. *Geoderma* 122, 1–23.
<https://doi.org/10.1016/j.geoderma.2004.01.021>
- Freudenschuß, A., 2010. Arbeiten zur Evaluierung von ÖPUL-Maßnahmen hinsichtlich ihrer Klimawirksamkeit: Schwerpunkt agrarische Bewirtschaftung. Umweltbundesamt.
- Gaiser, T., Stahr, K., Billen, N., Mohammad, M.A.-R., 2008. Modeling carbon sequestration under zero tillage at the regional scale. I. The effect of soil erosion. *Ecol. Model.* 218, 110–120.
- González-Sánchez, E.J., Ordóñez-Fernández, R., Carbonell-Bojollo, R., Veroz-González, O., Gil-Ribes, J.A., 2012. Meta-analysis on atmospheric carbon capture in Spain through the use of conservation agriculture. *Soil Tillage Res.* 122, 52–60.
<https://doi.org/10.1016/j.still.2012.03.001>
- Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., Bruni, E., Caliman, J.-P., Cardinael, R., Chen, S., Ciais, P., Desbois, D., Fouche, J., Frank, S., Henault, C., Lugato, E., Naipal, V., Nesme, T., Obersteiner, M., Pellerin, S., Powlson, D.S., Rasse, D.P., Rees, F., Soussana, J.-F., Su, Y., Tian, H., Valin, H., Zhou, F., 2021a. Can N₂O emissions offset the benefits from soil organic carbon storage? *Glob. Change Biol.* 27, 237–256.
<https://doi.org/10.1111/gcb.15342>
- Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., Bruni, E., Caliman, J.-P., Cardinael, R., Chen, S., Ciais, P., Desbois, D., Fouche, J., Frank, S., Henault, C., Lugato, E., Naipal, V., Nesme, T., Obersteiner, M., Pellerin, S., Powlson, D.S., Rasse, D.P., Rees, F., Soussana, J.-F., Su, Y., Tian, H., Valin, H., Zhou, F., 2021b. Can N₂O emissions offset the benefits from soil organic carbon storage? *Glob. Change Biol.* 27, 237–256.
<https://doi.org/10.1111/gcb.15342>
- Haddaway, N.R., Hedlund, K., Jackson, L.E., Kätterer, T., Lugato, E., Thomsen, I.K., Jørgensen, H.B., Isberg, P.-E., 2017. How does tillage intensity affect soil organic carbon? A systematic review. *Environ. Evid.* 6, 30. <https://doi.org/10.1186/s13750-017-0108-9>
- Heikkinen, J., Ketoja, E., Nuutinen, V., Regina, K., 2013. Declining trend of carbon in Finnish cropland soils in 1974–2009. *Glob. Change Biol.* 19, 1456–1469. <https://doi.org/10.1111/gcb.12137>
- Hermle, S., Anken, T., Leifeld, J., Weiskopf, P., 2008. The effect of the tillage system on soil organic carbon content under moist, cold-temperate conditions. *Soil Tillage Res.* 98, 94–105.
<https://doi.org/10.1016/j.still.2007.10.010>
- Iocola, I., Bassu, S., Farina, R., Antichi, D., Basso, B., Bindi, M., Dalla Marta, A., Danuso, F., Doro, L., Ferrise, R., Giglio, L., Ginaldi, F., Mazzoncini, M., Mula, L., Orsini, R., Corti, G., Pasqui, M., Seddaiu, G., Tomozeiu, R., Ventrella, D., Villani, G., Roggero, P.P., 2017. Can conservation tillage mitigate climate change impacts in Mediterranean cereal systems? A soil organic carbon assessment using long term experiments. *Eur. J. Agron.* 90, 96–107.
<https://doi.org/10.1016/j.eja.2017.07.011>
- Jokubauskaite, I., Karčauskienė, D., Slepėtiene, A., Repsiene, R., Amaleviciute, K., 2016. Effect of different fertilization modes on soil organic carbon sequestration in acid soils. *Acta Agric. Scand. Sect. B—Soil Plant Sci.* 66, 647–652.
- Joosten, H., Clarke, D., 2002. Wise use of mires and peatlands: background and principles including a framework for decision-making. International Peat Society ; International Mire Conservation Group, Jyväskylä] : [Greifswald.
- Karlsson, A.-B., Boström, B., Jernbäcker, E., Lundblad, M., Mjureke, D., 2020. Vägen till en klimatpositiv framtid. SOU 2020: Betänkande från Klimatpolitiska vägvalsutredningen (M 2018:07).



- Kauer, K., Astover, A., Viiralt, R., Raave, H., Kätterer, T., 2019. Evolution of soil organic carbon in a carbonaceous glacial till as an effect of crop and fertility management over 50 years in a field experiment. *Agric. Ecosyst. Environ.* 283, 106562.
- Keel, S.G., Anken, T., Büchi, L., Chervet, A., Fliessbach, A., Flisch, R., Huguenin-Elie, O., Mäder, P., Mayer, J., Sinaj, S., Sturny, W., Wüst-Galley, C., Zihlmann, U., Leifeld, J., 2019. Loss of soil organic carbon in Swiss long-term agricultural experiments over a wide range of management practices. *Agric. Ecosyst. Environ.* 286, 106654. <https://doi.org/10.1016/j.agee.2019.106654>
- Kögel-Knabner, I., Amelung, W., 2021. Soil organic matter in major pedogenic soil groups. *Geoderma* 384, 114785. <https://doi.org/10.1016/j.geoderma.2020.114785>
- Koopmans, C., Timmermans, B., Wagenaar, J.P., van, J., 2019. Evaluatie van maatregelen voor het vastleggen van koolstof 54.
- Kovats, R.S., Valentini, R., Bouwer, L.M., Georgopoulou, E., Jacob, D., Martin, E., Rounsevell, M., Soussana, J.-F., 2014. Europe, in: *Impacts, Adaptation, and Vulnerability. Part B: Regional Aspects. Contribution of Working Group II to the Fifth Assessment Report of the Intergovernmental Panel on Climate Change.* United Kingdom and New York.
- Lagomarsino, A., Agnelli, A.E., Linqvist, B., Adviento-Borbe, M.A., Agnelli, A., Gavina, G., Ravaglia, S., Ferrara, R.M., 2016. Alternate Wetting and Drying of Rice Reduced CH₄ Emissions but Triggered N₂O Peaks in a Clayey Soil of Central Italy. *Pedosphere* 26, 533–548. [https://doi.org/10.1016/S1002-0160\(15\)60063-7](https://doi.org/10.1016/S1002-0160(15)60063-7)
- Lagomarsino, A., Agnelli, A.E., Linqvist, B., Adviento-borbe, M.A., Agnelli, A., Gavina, G., Ravaglia, S., Ferrara, R.M., 2016. Alternate Wetting and Drying of Rice Reduced CH₄ Emissions but Triggered N₂O Peaks in a Clayey Soil of Central Italy. *Pedosphere* 26, 533–548. [https://doi.org/10.1016/S1002-0160\(15\)60063-7](https://doi.org/10.1016/S1002-0160(15)60063-7)
- Lal, R., 2018. Digging deeper: A holistic perspective of factors affecting soil organic carbon sequestration in agroecosystems. *Glob. Change Biol.* 24, 3285–3301. <https://doi.org/10.1111/gcb.14054>
- Lal, R., 2004. Soil carbon sequestration to mitigate climate change. *Geoderma* 123, 1–22. <https://doi.org/10.1016/j.geoderma.2004.01.032>
- Lanigan, G., Donnellan, T., Hanrahan, K., Carsten, P., Shalloo, L., Krol, D., Forrester, P.J., Farrelly, N., O'Brien, D., Ryan, M., Murphy, P., Caslin, B., Spink, J., Finnan, J., Boland, A., Upton, J., Richards, K.G., 2018. *An Analysis of Abatement Potential of Greenhouse Gas Emissions in Irish Agriculture 2021-2030 (Technical Report).* Teagasc.
- Lanigan, G., Forristal, D., Prescher, A., 2012. *Assesing the greenhouse gas budget of tillage mitigation options for arable systems.* Technology updates, Crops environment and Land Use. Agriculture and Food development authority, Teagasc. h.
- Leifeld, J., Bassin, S., Fuhrer, J., 2003. Carbon stocks and carbon sequestration potentials in agricultural soils in Switzerland. *Agroscope, FAL-Schriftenreihe 44*, Zürich. - Google Search (Schriftreihe FAL No. 44). Eidgenössische Forschungsanstalt für Agrarökologie und Landbau, FAL Reckenholz, Zürich.
- Leifeld, J., Menichetti, L., 2018. The underappreciated potential of peatlands in global climate change mitigation strategies. *Nat. Commun.* 9, 1071. <https://doi.org/10.1038/s41467-018-03406-6>
- Lesschen, J.P., Heesmans, H.I.M., Mol-Dijkstra, J.P., Doorn, A.M. van, Verkaik, E., Wyngaert, I.J.J. van den, Kuikman, P.J., 2012. *Mogelijkheden voor koolstofvastlegging in de Nederlandse landbouw en natuur.* Alterra.
- López-Bellido, L., López-Bellido, R., Fernández-García, P., Muñoz-Romero, V., Lopez-Bellido, F.J., 2020. Carbon storage in a rainfed Mediterranean vertisol: Effects of tillage and crop rotation in a long-term experiment. *Eur. J. Soil Sci.* 71, 472–483. <https://doi.org/10.1111/ejss.12883>



- Lugato, E., Bampa, F., Panagos, P., Montanarella, L., Jones, A., 2014. Potential carbon sequestration of European arable soils estimated by modelling a comprehensive set of management practices. *Glob. Change Biol.* 20, 3557–3567. <https://doi.org/10.1111/gcb.12551>
- Lugato, E., Berti, A., Giardini, L., 2006. Soil organic carbon (SOC) dynamics with and without residue incorporation in relation to different nitrogen fertilisation rates. *Geoderma* 135, 315–321. <https://doi.org/10.1016/j.geoderma.2006.01.012>
- Lugato, E., Leip, A., Jones, A., 2018. Mitigation potential of soil carbon management overestimated by neglecting N₂O emissions. *Nat. Clim. Change* 8, 219–223. <https://doi.org/10.1038/s41558-018-0087-z>
- MacLeod, M., Moran, D., Eory, V., Rees, R.M., Barnes, A., Topp, C.F.E., Ball, B., Hoad, S., Wall, E., McVittie, A., Pajot, G., Matthews, R., Smith, P., Moxey, A., 2010. Developing greenhouse gas marginal abatement cost curves for agricultural emissions from crops and soils in the UK. *Agric. Syst.* 103, 198–209. <https://doi.org/10.1016/j.agsy.2010.01.002>
- Martínez, I., Chervet, A., Weisskopf, P., Sturny, W.G., Etana, A., Stettler, M., Forkman, J., Keller, T., 2016. Two decades of no-till in the Oberacker long-term field experiment: Part I. Crop yield, soil organic carbon and nutrient distribution in the soil profile. *Soil Tillage Res.* 163, 141–151. <https://doi.org/10.1016/j.still.2016.05.021>
- Mazzoncini, M., Antichi, D., Di Bene, C., Risaliti, R., Petri, M., Bonari, E., 2016. Soil carbon and nitrogen changes after 28 years of no-tillage management under Mediterranean conditions. *Eur. J. Agron.* 77, 156–165. <https://doi.org/10.1016/j.eja.2016.02.011>
- Mazzoncini, M., Sapkota, T.B., Bàrberi, P., Antichi, D., Risaliti, R., 2011. Long-term effect of tillage, nitrogen fertilization and cover crops on soil organic carbon and total nitrogen content. *Soil Tillage Res.* 114, 165–174. <https://doi.org/10.1016/j.still.2011.05.001>
- Meersmans, J., Arrouays, D., Van Rompaey, A.J.J., Pagé, C., De Baets, S., Quine, T.A., 2016. Future C loss in mid-latitude mineral soils: climate change exceeds land use mitigation potential in France. *Sci. Rep.* 6, 35798. <https://doi.org/10.1038/srep35798>
- Meijide, A., Gruening, C., Goded, I., Seufert, G., Cescatti, A., 2017. Water management reduces greenhouse gas emissions in a Mediterranean rice paddy field. *Quantif. Mitig. Greenh. Gas Emiss. Mediterr. Crop. Syst.* 238, 168–178. <https://doi.org/10.1016/j.agee.2016.08.017>
- Metzger, M.J., Bunce, R.G.H., Jongman, R.H.G., Mùcher, C.A., Watkins, J.W., 2005. A climatic stratification of the environment of Europe. *Glob. Ecol. Biogeogr.* 14, 549–563. <https://doi.org/10.1111/j.1466-822X.2005.00190.x>
- Meurer, K., Barron, J., Chenu, C., Coucheney, E., Fielding, M., Hallett, P., Herrmann, A.M., Keller, T., Koestel, J., Larsbo, M., Lewan, E., Or, D., Parsons, D., Parvin, N., Taylor, A., Vereecken, H., Jarvis, N., 2020. A framework for modelling soil structure dynamics induced by biological activity. *Glob. Change Biol.* 26, 5382–5403. <https://doi.org/10.1111/gcb.15289>
- Meurer, K.H., Haddaway, N.R., Bolinder, M.A., Kätterer, T., 2018. Tillage intensity affects total SOC stocks in boreo-temperate regions only in the topsoil—A systematic review using an ESM approach. *Earth-Sci. Rev.* 177, 613–622.
- Mondini, C., Coleman, K., Whitmore, A.P., 2012. Spatially explicit modelling of changes in soil organic C in agricultural soils in Italy, 2001–2100: Potential for compost amendment. *Agric. Ecosyst. Environ.* 153, 24–32. <https://doi.org/10.1016/j.agee.2012.02.020>
- Moreno-García, M., Repullo-Ruibérriz de Torres, M.Á., González-Sánchez, E.J., Ordóñez-Fernández, R., Veroz-González, Ó., Carbonell-Bojollo, R.M., 2020. Methodology for estimating the impact of no tillage on the 4perMille initiative: The case of annual crops in Spain. *Geoderma* 371, 114381. <https://doi.org/10.1016/j.geoderma.2020.114381>
- Nabuurs, G., Wp, D., Aj, D., O, O., E, V., P, K., Ap, W., Gmj, M., 1999. Resolving Issues on Terrestrial Biospheric Sinks in the Kyoto Protocol. Het oplossen van problemen rond de aardse biosferische koolstof verzinkingen in de Kyoto protocol.



- NIR 2020 [WWW Document], n.d. URL <http://www.sinanet.isprambiente.it/it/sia-ispra/serie-storiche-emissioni/national-inventory-report/view> (accessed 1.19.21).
- Olson, K.R., Al-Kaisi, M.M., Lal, R., Lowery, B., 2014. Experimental Consideration, Treatments, and Methods in Determining Soil Organic Carbon Sequestration Rates. *Soil Sci. Soc. Am. J.* 78, 348–360. <https://doi.org/10.2136/sssaj2013.09.0412>
- Paul, C., Fealy, R., Fenton, O., Lanigan, G., O’Sullivan, L., Schulte, R.P.O., 2018. Assessing the role of artificially drained agricultural land for climate change mitigation in Ireland. *Environ. Sci. Policy* 80, 95–104. <https://doi.org/10.1016/j.envsci.2017.11.004>
- Paustian, K., Lehmann, J., Ogle, S., Reay, D., Robertson, G.P., Smith, P., 2016. Climate-smart soils. *Nature* 532, 49–57. <https://doi.org/10.1038/nature17174>
- Pellerin, S., Bamière, L., Angers, D., Béline, F., Benoît, M., Butault, J.P., 2013. How can French agriculture contribute to reducing greenhouse gas emissions? *Energy* 19, 12.
- Pellerin, S., Bamière, L., Launay, C., Martin, R., Schiavo, M., Angers, D., Augusto, L., Balesdent, J., Doelsch, I.B., Bellassen, V., 2019. Stocker du carbone dans les sols français, quel potentiel au regard de l’objectif 4 pour 1000 et à quel coût?
- Peyron, M., Bertora, C., Pelissetti, S., Said-Pullicino, D., Celi, L., Miniotti, E., Romani, M., Sacco, D., 2016. Greenhouse gas emissions as affected by different water management practices in temperate rice paddies. *Agric. Ecosyst. Environ.* 232, 17–28. <https://doi.org/10.1016/j.agee.2016.07.021>
- Poeplau, C., Don, A., 2015. Carbon sequestration in agricultural soils via cultivation of cover crops – A meta-analysis. *Agric. Ecosyst. Environ.* 200, 33–41. <https://doi.org/10.1016/j.agee.2014.10.024>
- Poeplau, C., Kätterer, T., Bolinder, M.A., Börjesson, G., Berti, A., Lugato, E., 2015. Low stabilization of aboveground crop residue carbon in sandy soils of Swedish long-term experiments. *Geoderma* 237, 246–255.
- Poeplau, C., Zopf, D., Greiner, B., Geerts, R., Korvaar, H., Thumm, U., Don, A., Heidkamp, A., Flessa, H., 2018. Why does mineral fertilization increase soil carbon stocks in temperate grasslands? *Agric. Ecosyst. Environ.* 265, 144–155.
- Poulton, P., Johnston, J., Macdonald, A., White, R., Powlson, D., 2018a. Major limitations to achieving “4 per 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term experiments at Rothamsted Research, United Kingdom. *Glob. Change Biol.* 24, 2563–2584.
- Poulton, P., Johnston, J., Macdonald, A., White, R., Powlson, D., 2018b. Major limitations to achieving “4 per 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term experiments at Rothamsted Research, United Kingdom. *Glob. Change Biol.* 24, 2563–2584. <https://doi.org/10.1111/gcb.14066>
- Powlson, D.S., Bhogal, A., Chambers, B.J., Coleman, K., Macdonald, A.J., Goulding, K.W.T., Whitmore, A.P., 2012. The potential to increase soil carbon stocks through reduced tillage or organic material additions in England and Wales: a case study. *Agric. Ecosyst. Environ.* 146, 23–33.
- Powlson, D.S., Whitmore, A.P., Goulding, K.W.T., 2011. Soil carbon sequestration to mitigate climate change: a critical re-examination to identify the true and the false. *Eur. J. Soil Sci.* 62, 42–55. <https://doi.org/10.1111/j.1365-2389.2010.01342.x>
- Rasse, D., Økland, I.H., Bárcena, T.G., Riley, H., Martinsen, V., Sturite, I., Jøner, E., O’Toole, A., Øpstad, S., Cottis, T., 2019. Muligheter og utfordringer for økt karbonbinding i jordbruksjord. NIBIO Rapp.
- Reay, D.S., Davidson, E.A., Smith, K.A., Smith, P., Melillo, J.M., Dentener, F., Crutzen, P.J., 2012. Global agriculture and nitrous oxide emissions. *Nat. Clim. Change* 2, 410–416. <https://doi.org/10.1038/nclimate1458>
- Renou-Wilson, F., Barry, C., Müller, C., Wilson, D., 2014. The impacts of drainage, nutrient status and management practice on the full carbon balance of grasslands on organic soils in a maritime



- temperate zone. *Biogeosciences Discuss.* 11, 5557–5612. <https://doi.org/10.5194/bgdc-11-5557-2014>
- Roelandt, C., Dendoncker, N., Rounsevell, M., Perrin, D., Wesemael, B.V., 2007. Projecting future N₂O emissions from agricultural soils in Belgium. *Glob. Change Biol.* 13, 18–27. <https://doi.org/10.1111/j.1365-2486.2006.01273.x>
- Roelandt, C., Wesemael, B.V., Rounsevell, M., 2005. Estimating annual N₂O emissions from agricultural soils in temperate climates. *Glob. Change Biol.* 11, 1701–1711. <https://doi.org/10.1111/j.1365-2486.2005.01025.x>
- Sanderman, J., Hengl, T., Fiske, G.J., 2017. Soil carbon debt of 12,000 years of human land use. *Proc. Natl. Acad. Sci.* 114, 9575–9580. <https://doi.org/10.1073/pnas.1706103114>
- Schlesinger, W.H., 2000. Carbon sequestration in soils: some cautions amidst optimism. *Agric. Ecosyst. Environ.* 82, 121–127. [https://doi.org/10.1016/S0167-8809\(00\)00221-8](https://doi.org/10.1016/S0167-8809(00)00221-8)
- Schneider, F., Don, A., 2019. Root-restricting layers in German agricultural soils. Part II: Adaptation and melioration strategies. *Plant Soil* 442, 419–432. <https://doi.org/10.1007/s11104-019-04186-8>
- Simo, I., Schulte, R., O’Sullivan, L., Creamer, R., 2019. Digging deeper: Understanding the contribution of subsoil carbon for climate mitigation, a case study of Ireland. *Environ. Sci. Policy* 98, 61–69. <https://doi.org/10.1016/j.envsci.2019.05.004>
- Singh, B.R., Lal, R., 2005. The potential of soil carbon sequestration through improved management practices in Norway. *Environ. Dev. Sustain.* 7, 161–184.
- Sinsabaugh, R.L., Manzoni, S., Moorhead, D.L., Richter, A., 2013. Carbon use efficiency of microbial communities: stoichiometry, methodology and modelling. *Ecol. Lett.* 16, 930–939. <https://doi.org/10.1111/ele.12113>
- Sleutel, S., De Neve, S., Hofman, G., 2007. Assessing causes of recent organic carbon losses from cropland soils by means of regional-scaled input balances for the case of Flanders (Belgium). *Nutr. Cycl. Agroecosystems* 78, 265–278.
- Smith, P., 2014. Do grasslands act as a perpetual sink for carbon? *Glob. Change Biol.* 20, 2708–2711. <https://doi.org/10.1111/gcb.12561>
- Smith, P., Falloon, P., Griffiths, H., Jarvis, P., 2005. Carbon sequestration in European croplands. *Carbon Balance For. Biomes* 47–55.
- Smith, P., Martino, D., Cai, Z., Gwary, D., Janzen, H., Kumar, P., McCarl, B., Ogle, S., O’Mara, F., Rice, C., Scholes, B., Sirotenko, O., 2007. Agriculture, in: *In Climate Change: Mitigation. Contribution of Working Group III to the Fourth Assessment Report of the Intergovernmental Panel on Climate Change.* Cambridge University Press, Cambridge, United Kingdom and New York, NY, USA, pp. 498–540.
- Smith, P., Powlson, D.S., Glendining, M.J., Smith, J.U., 1998. Preliminary estimates of the potential for carbon mitigation in European soils through no-till farming. *Glob. Change Biol.* 4, 679–685. <https://doi.org/10.1046/j.1365-2486.1998.00185.x>
- Smith, P., Powlson, D.S., Glendining, M.J., Smith, J.U., 1996. Opportunities and limitations for C sequestration in European agricultural soils through changes in management.
- Smith, P., Powlson, D.S., Smith, J.U., Falloon, P., Coleman, K., 2000. Meeting Europe’s climate change commitments: quantitative estimates of the potential for carbon mitigation by agriculture. *Glob. Change Biol.* 6, 525–539. <https://doi.org/10.1046/j.1365-2486.2000.00331.x>
- Smith, P., Soussana, J.-F., Angers, D., Schipper, L., Chenu, C., Rasse, D.P., Batjes, N.H., Egmond, F. van, McNeill, S., Kuhnert, M., Arias-Navarro, C., Olesen, J.E., Chirinda, N., Fornara, D., Wollenberg, E., Álvaro-Fuentes, J., Sanz-Cobena, A., Klumpp, K., 2020. How to measure, report and verify soil carbon change to realize the potential of soil carbon sequestration for atmospheric greenhouse gas removal. *Glob. Change Biol.* 26, 219–241. <https://doi.org/10.1111/gcb.14815>



- Smith, W.N., Desjardins, R.L., Pattey, E., 2000. The net flux of carbon from agricultural soils in Canada 1970–2010. *Glob. Change Biol.* 6, 557–568. <https://doi.org/10.1046/j.1365-2486.2000.00340.x>
- Stockmann, U., Adams, M.A., Crawford, J.W., Field, D.J., Henakaarchchi, N., Jenkins, M., Minasny, B., McBratney, A.B., Courcelles, V. de R. de, Singh, K., Wheeler, I., Abbott, L., Angers, D.A., Baldock, J., Bird, M., Brookes, P.C., Chenu, C., Jastrow, J.D., Lal, R., Lehmann, J., O'Donnell, A.G., Parton, W.J., Whitehead, D., Zimmermann, M., 2013. The knowns, known unknowns and unknowns of sequestration of soil organic carbon. *Agric. Ecosyst. Environ.* 164, 80–99. <https://doi.org/10.1016/j.agee.2012.10.001>
- Taghizadeh-Toosi, A., Christensen, B.T., Hutchings, N.J., Vejlin, J., Kätterer, T., Glendining, M., Olesen, J.E., 2014. C-TOOL: A simple model for simulating whole-profile carbon storage in temperate agricultural soils. *Ecol. Model.* 292, 11–25. <https://doi.org/10.1016/j.ecolmodel.2014.08.016>
- Taghizadeh-Toosi, A., Olesen, J.E., 2016. Modelling soil organic carbon in Danish agricultural soils suggests low potential for future carbon sequestration. *Agric. Syst.* 145, 83–89.
- Taghizadeh-Toosi, A., Olesen, J.E., Kristensen, K., Elsgaard, L., Østergaard, H.S., Lægdsmand, M., Greve, M.H., Christensen, B.T., 2014. Changes in carbon stocks of Danish agricultural mineral soils between 1986 and 2009. *Eur. J. Soil Sci.* 65, 730–740.
- Tanneberger, F., Moen, A., Joosten, H., Nilsen, N., 2017. The peatland map of Europe.
- Teixeira, R.F.M., Domingos, T., Costa, A., Oliveira, R., Farropas, L., Calouro, F., Barradas, A.M., Carneiro, J., 2011. Soil organic matter dynamics in Portuguese natural and sown rainfed grasslands. *Ecol. Model.* 222, 993–1001.
- Tiemeyer, B., Albiac Borraz, E., Augustin, J., Bechtold, M., Beetz, S., Beyer, C., Drösler, M., Ebli, M., Eickenscheidt, T., Fiedler, S., 2016. High emissions of greenhouse gases from grasslands on peat and other organic soils. *Glob. Change Biol.* 22, 4134–4149.
- Torres-Sallan, G., Schulte, R., Lanigan, G.J., Byrne, K.A., Reidy, B., Simo, I., Six, J., Creamer, R.E., 2017. Clay illuviation provides a longterm sink for C sequestration in subsoils. <https://doi.org/10.1038/srep45635>
- Triberti, L., Nastri, A., Giordani, G., Comellini, F., Baldoni, G., Toderi, G., 2008. Can mineral and organic fertilization help sequester carbon dioxide in cropland? *Eur. J. Agron.* 29, 13–20. <https://doi.org/10.1016/j.eja.2008.01.009>
- van Groenigen, K.J., Hastings, A., Forristal, D., Roth, B., Jones, M., Smith, P., 2011. Soil C storage as affected by tillage and straw management: An assessment using field measurements and model predictions. *Agric. Ecosyst. Environ.* 140, 218–225. <https://doi.org/10.1016/j.agee.2010.12.008>
- Vanden Nest, T., Ruyschaert, G., Vandecasteele, B., Cougnon, M., Merckx, R., Reheul, D., 2015. P availability and P leaching after reducing the mineral P fertilization and the use of digestate products as new organic fertilizers in a 4-year field trial with high P status. *Agric. Ecosyst. Environ.* 202, 56–67. <https://doi.org/10.1016/j.agee.2014.12.012>
- Vleeshouwers, L.M., Verhagen, A., 2002. Carbon emission and sequestration by agricultural land use: a model study for Europe. *Glob. Change Biol.* 8, 519–530. <https://doi.org/10.1046/j.1365-2486.2002.00485.x>
- Wiesmeier, M., Mayer, S., Burmeister, J., Hübner, R., Kögel-Knabner, I., 2020. Feasibility of the 4 per 1000 initiative in Bavaria: A reality check of agricultural soil management and carbon sequestration scenarios. *Geoderma* 369, 114333. <https://doi.org/10.1016/j.geoderma.2020.114333>
- Wikström, L., 2019. Så klarar det svenska jordbruket klimatmålen: en delrapport från IVA-projektet Vägval för klimatet.
- Willekens, K., Vandecasteele, B., Buchan, D., De Neve, S., 2014. Soil quality is positively affected by reduced tillage and compost in an intensive vegetable cropping system. *Appl. Soil Ecol.* 82, 61–71. <https://doi.org/10.1016/j.apsoil.2014.05.009>



- Wilson, D., Blain, D., Couwenberg, J., Evans, C.D., Murdiyarso, D., Page, S.E., Renou-Wilson, F., Rieley, J.O., Sirin, A., Strack, M., 2016. Greenhouse gas emission factors associated with rewetting of organic soils. *Mires Peat* 17.
- Wüst-Galley, C., Grünig, A., Leifeld, J., 2020. Land use-driven historical soil carbon losses in Swiss peatlands. *Landsc. Ecol.* 35, 173–187. <https://doi.org/10.1007/s10980-019-00941-5>

Annex

A List of references mineral soils

Austria:

- Chen, J., Heiling, M., Resch, C., Mbaye, M., Gruber, R., Dercon, G., 2018. Does maize and legume crop residue mulch matter in soil organic carbon sequestration? *Agriculture, ecosystems & environment* 265, 123–131.
- Dersch, G., Böhm, K., 2001. Effects of agronomic practices on the soil carbon storage potential in arable farming in Austria. *Nutrient Cycling in Agroecosystems* 60, 49–55.
- Freudenschuß, A., 2010. Arbeiten zur Evaluierung von ÖPUL-Maßnahmen hinsichtlich ihrer Klimawirksamkeit: Schwerpunkt agrarische Bewirtschaftung. Umweltbundesamt.
- Kasper, M., Freyer, B., Hülsbergen, K.-J., Schmid, H., Friedel, J.K., 2015. Humus balances of different farm production systems in main production areas in Austria. *Journal of Plant Nutrition and Soil Science* 178, 25–34.
- Sedy, K., 2015. Project ACC -Austrian Carbon Calculator. ACRP 4th Call. (Final report - Activity report).
- Spiegel, H., Dersch, G., Hösch, J., Baumgarten, A., 2007. Tillage effects on soil organic carbon and nutrient availability in a long-term field experiment in Austria. *Die Bodenkultur* 58, 47–58.

Belgium:

Flanders

- Debode, J., Ebrahimi, N., D'Hose, T., Cremelie, P., Viaene, N., Vandecasteele, B., 2020. Has compost with biochar added during the process added value over biochar or compost to increase disease suppression? *Applied Soil Ecology* 153, 103571. <https://doi.org/10.1016/j.apsoil.2020.103571>
- D'Haene, K., 2008. The potential of reduced tillage agriculture in Flanders. s.n., S.l.
- D'Hose, T., Ruysschaert, G., 2017. Mogelijkheden voor koolstofopslag onder gras- en akkerland in Vlaanderen. (No. No. 231), ILVO Mededeling. Netherlands.
- D'Hose, T., Ruysschaert, G., Viaene, N., Debode, J., Vanden Nest, T., Van Vaerenbergh, J., Cornelis, W., Willekens, K., Vandecasteele, B., 2016. Farm compost amendment and non-inversion tillage improve soil quality without increasing the risk for N and P leaching. *Agriculture, Ecosystems & Environment* 225, 126–139. <https://doi.org/10.1016/j.agee.2016.03.035>
- Mestdagh, I., Lootens, P., Van Cleemput, O., Carlier, L., 2006. Variation in organic-carbon concentration and bulk density in Flemish grassland soils. *Journal of plant nutrition and soil science* 169, 616–622.
- Mestdagh, I.P.M., 2005. Carbon sequestration in different Flemish grassland ecosystems.
- Nelissen, V., Ruysschaert, G., Manka'Abusi, D., D'Hose, T., De Beuf, K., Al-Barri, B., Cornelis, W., Boeckx, P., 2015. Impact of a woody biochar on properties of a sandy loam soil and spring barley during a two-year field experiment. *European Journal of Agronomy* 62, 65–78. <https://doi.org/10.1016/j.eja.2014.09.006>



- Nest, T.V., Ruyschaert, G., Vandecasteele, B., Cougnon, M., Merckx, R., Reheul, D., 2015. P availability and P leaching after reducing the mineral P fertilization and the use of digestate products as new organic fertilizers in a 4-year field trial with high P status. *Agriculture, Ecosystems & Environment* 202, 56–67. <https://doi.org/10.1016/j.agee.2014.12.012>
- Pardon, P., Reubens, B., Reheul, D., Mertens, J., De Frenne, P., Coussement, T., Janssens, P., Verheyen, K., 2017. Trees increase soil organic carbon and nutrient availability in temperate agroforestry systems. *Agriculture, Ecosystems & Environment* 247, 98–111. <https://doi.org/10.1016/j.agee.2017.06.018>
- Sleutel, S., De Neve, S., Hofman, G., 2007. Assessing causes of recent organic carbon losses from cropland soils by means of regional-scaled input balances for the case of Flanders (Belgium). *Nutrient Cycling in Agroecosystems* 78, 265–278.
- Van Vooren, L., 2018. Multifunctionality in agriculture: impact of hedgerows, grass strips and extensive grassland management on crops, regulating ecosystem services and biodiversity. Ghent University.
- Vermang, J., 2012. Erosion processes and physical quality of loamy soils as affected by reduced tillage. PhD thesis. Ghent University, Belgium (PhD-thesis). Ghent University, Belgium.
- Viaene, J., Van Lancker, J., Vandecasteele, B., Willekens, K., Bijttebier, J., Ruyschaert, G., De Neve, S., Reubens, B., 2016. Opportunities and barriers to on-farm composting and compost application: A case study from northwestern Europe. *Waste Management* 48, 181–192.
- Willekens, K., Vandecasteele, B., Buchan, D., De Neve, S., 2014. Soil quality is positively affected by reduced tillage and compost in an intensive vegetable cropping system. *Applied Soil Ecology* 82, 61–71. <https://doi.org/10.1016/j.apsoil.2014.05.009>
- Xu, H., Vandecasteele, B., Zavattaro, L., Sacco, D., Wendland, M., Boeckx, P., Haesaert, G., Sleutel, S., 2019. Maize root-derived C in soil and the role of physical protection on its relative stability over shoot-derived C. *European Journal of Soil Science* 70, 935–946.

Wallonia

- Buysse, P., Roisin, C., Aubinet, M., 2013. Fifty years of contrasted residue management of an agricultural crop: Impacts on the soil carbon budget and on soil heterotrophic respiration. *Agriculture, Ecosystems & Environment* 167, 52–59. <https://doi.org/10.1016/j.agee.2013.01.006>
- Chartin, C., Carnol, M., van Wesemael, B., 2018. CARBIOSOL 4-Développement d'indicateurs de la qualité biologique et du carbone organique du sol pour l'évaluation de l'état des sols en Wallonie. ULiège.
- Chartin, C., et al., Mapping Soil Organic Carbon stocks and estimating uncertainties at the regional scale following a legacy sampling strategy (Southern Belgium, Wallonia), <http://dx.doi.org/10.1016/j.geodrs.2016.12.006>
- Dendoncker, N., Van Wesemael, B., Rounsevell, M.D.A., Roelandt, C., Lettens, S., 2004. Belgium's CO₂ mitigation potential under improved cropland management. *Agriculture, Ecosystems & Environment* 103, 101–116. <https://doi.org/10.1016/j.agee.2003.10.010>
- Goidts, E., van Wesemael, B., 2007. Regional assessment of soil organic carbon changes under agriculture in Southern Belgium (1955–2005). *Geoderma* 141, 341–354. <https://doi.org/10.1016/j.geoderma.2007.06.013>
- Gourlez de la Motte, L., 2019. Carbon balance of an intensively managed pasture: methodology, evaluation and impacts of weather conditions and grazing strategy. Université de Liège, Liège, Belgique.
- Lognoul, M., Debacq, A., De Ligne, A., Dumont, B., Manise, T., Bodson, B., Heinesch, B., Aubinet, M., 2019. N₂O flux short-term response to temperature and topsoil disturbance in a fertilized crop: An eddy covariance campaign. *Agricultural and Forest Meteorology* 271, 193–206. <https://doi.org/10.1016/j.agrformet.2019.02.033>



Lognoul, M., Theodorakopoulos, N., Hiel, M.-P., Regaert, D., Broux, F., Heinesch, B., Bodson, B., Vandenbol, M., Aubinet, M., 2017. Impact of tillage on greenhouse gas emissions by an agricultural crop and dynamics of N₂O fluxes: Insights from automated closed chamber measurements. *Soil and Tillage Research* 167, 80–89.
<https://doi.org/10.1016/j.still.2016.11.008>

Van Orshoven, J., Maes, J., Vereecken, H., Feyen, J., Dudal, R., 1988. A structured database of Belgian soil profile data. *Pedologie: bulletin van de Belgische bodemkundige vereniging* 38, 191–206.

Czech Republic

Šimon, T., Javůrek, M., Mikanová, O., Vach, M. 2009: The influence of tillage systems on soil organic matter and soil hydrophobicity. *Soil and Tillage Research* 105, 44–48.

Denmark

Taghizadeh-Toosi, A., Olesen, J.E., 2016. Modelling soil organic carbon in Danish agricultural soils suggests low potential for future carbon sequestration. *Agricultural Systems* 145, 83–89.
<https://doi.org/10.1016/j.agsy.2016.03.004>

Taghizadeh-Toosi, A., Christensen, B.T., Hutchings, N.J., Vejlin, J., Kätterer, T., Glendining, M., Olesen, J.E., 2014. C-TOOL: A simple model for simulating whole-profile carbon storage in temperate agricultural soils. *Ecological Modelling* 292, 11–25.
<https://doi.org/10.1016/j.ecolmodel.2014.08.016>

Taghizadeh-Toosi, A., Olesen, J.E., Kristensen, K., Elsgaard, L., Østergaard, H.S., Lægdsmand, M., Greve, M.H., Christensen, B.T., 2014a. Changes in carbon stocks of Danish agricultural mineral soils between 1986 and 2009. *European Journal of Soil Science* 65, 730–740.

Estonia

Kauer, K., Astover, A., Viiralt, R., Raave, H., Kätterer, T., 2019. Evolution of soil organic carbon in a carbonaceous glacial till as an effect of crop and fertility management over 50 years in a field experiment. *Agriculture, Ecosystems & Environment* 283, 106562.

Kauer, K., Kölli, R., Viiralt, R., Köster, T., Noormets, M., Laidna, T., Keres, I., Parol, A., Varul, T., Selge, A., 2013. Effect of cut plant residue management and fertilization on the dry-matter yield of swards and on carbon content of soil. *Communications in soil science and plant analysis* 44, 205–218.

Kauer, K., Tein, B., de Cima, D.S., Talgre, L., Eremeev, V., Loit, E., Luik, A., 2015. Soil carbon dynamics estimation and dependence on farming system in a temperate climate. *Soil and Tillage Research* 154, 53–63.

Finland

Singh P., Heikkinen, J., Ketoje, E., Nuutinen, V., Palojärvi, A., Sheehy, J., Esala, M., Mitra, S., Alakukku, L. and Regina, K. (2015) Tillage and crop residue management methods had minor effects on the stock and stabilization of topsoil carbon in a 30-year field experiment. *Science of the Total Environment* 518–519 (2015) 337–344.

France

Autret, B., Mary, B., Chenu, C., Balabane, M., Girardin, C., Bertrand, M., Grandeau, G., Beudoin, N., 2016. Alternative arable cropping systems: A key to increase soil organic carbon storage? Results from a 16 year field experiment. *Agriculture, Ecosystems & Environment* 232, 150–164. <https://doi.org/10.1016/j.agee.2016.07.008>

Cardinael, R., Chevallier, T., Barthès, B.G., Saby, N.P.A., Parent, T., Dupraz, C., Bernoux, M., Chenu, C., 2015. Impact of alley cropping agroforestry on stocks, forms and spatial distribution of soil organic carbon — A case study in a Mediterranean context. *Geoderma* 259–260, 288–299.



- <https://doi.org/10.1016/j.geoderma.2015.06.015>
- Cardinael, R., Chevallier, T., Cambou, A., Béral, C., Barthès, B.G., Dupraz, C., Durand, C., Kouakoua, E., Chenu, C., 2017. Increased soil organic carbon stocks under agroforestry: A survey of six different sites in France. *Agriculture, Ecosystems & Environment* 236, 243–255. <https://doi.org/10.1016/j.agee.2016.12.011>
- Chen, S., Arrouays, D., Angers, D.A., Chenu, C., Barré, P., Martin, M.P., Saby, N.P.A., Walter, C., 2019. National estimation of soil organic carbon storage potential for arable soils: A data-driven approach coupled with carbon-landscape zones. *Science of The Total Environment* 666, 355–367. <https://doi.org/10.1016/j.scitotenv.2019.02.249>
- Chen, S., Martin, M.P., Saby, N.P.A., Walter, C., Angers, D.A., Arrouays, D., 2018. Fine resolution map of top- and subsoil carbon sequestration potential in France. *Science of The Total Environment* 630, 389–400. <https://doi.org/10.1016/j.scitotenv.2018.02.209>
- Constantin, J., Mary, B., Laurent, F., Aubrion, G., Fontaine, A., Kerveillant, P., Beaudoin, N., 2010. Effects of catch crops, no till and reduced nitrogen fertilization on nitrogen leaching and balance in three long-term experiments. *Agriculture, Ecosystems & Environment* 135, 268–278. <https://doi.org/10.1016/j.agee.2009.10.005>
- Dimassi, B., Cohan, J.-P., Labreuche, J., Mary, B., 2013. Changes in soil carbon and nitrogen following tillage conversion in a long-term experiment in Northern France. *Agriculture, Ecosystems & Environment* 169, 12–20. <https://doi.org/10.1016/j.agee.2013.01.012>
- Dimassi, B., Mary, B., Wylleman, R., Labreuche, J., Couture, D., Piraux, F., Cohan, J.-P., 2014. Long-term effect of contrasted tillage and crop management on soil carbon dynamics during 41 years. *Agriculture, Ecosystems & Environment* 188, 134–146. <https://doi.org/10.1016/j.agee.2014.02.014>
- Follain, S., Walter, C., Legout, A., Lemerrier, B., Dutin, G., 2007. Induced effects of hedgerow networks on soil organic carbon storage within an agricultural landscape. *Geoderma* 142, 80–95. <https://doi.org/10.1016/j.geoderma.2007.08.002>
- Guenet, B., Gabrielle, B., Chenu, C., Arrouays, D., Balesdent, J., Bernoux, M., Bruni, E., Caliman, J.-P., Cardinael, R., Chen, S., Ciais, P., Desbois, D., Fouche, J., Frank, S., Henault, C., Lugato, E., Naipal, V., Nesme, T., Obersteiner, M., Pellerin, S., Powlson, D.S., Rasse, D.P., Rees, F., Soussana, J.-F., Su, Y., Tian, H., Valin, H., Zhou, F., 2021. Can N₂O emissions offset the benefits from soil organic carbon storage? *Global Change Biology* 27, 237–256. <https://doi.org/10.1111/gcb.15342>
- Houot, S., Pons, M., Pradel, M., Tibi, A., Aubry, C., Augusto, L., Barbier, R., Benoit, P., Brugère, H., Caillaud, M.-A., Casellas, M., Chatelet, A., Dabert, P., De Marschal, S., 2014. Valorisation des matières fertilisantes d'origine résiduaire sur les sols à usage agricole ou forestier (Rapport final de l'expertise scientifique collective. Paris: INRA-CNRS-Irstea).
- Launay, C., Constantin, J., Chlebowski, F., Houot, S., Graux, A.-I., Klumpp, K., Martin, R., Mary, B., Pellerin, S., Therond, O., 2021 (accepted). Estimating the carbon storage potential and greenhouse gas emissions of French arable cropland using high-resolution modeling. *Global Change Biology* n/a. <https://doi.org/10.1111/gcb.15512>
- Levassasseur, F., Mary, B., Christensen, B.T., Duparque, A., Ferchaud, F., Kätterer, T., Lagrange, H., Montenach, D., Resseguier, C., Houot, S., 2020. The simple AMG model accurately simulates organic carbon storage in soils after repeated application of exogenous organic matter. *Nutr Cycl Agroecosyst* 117, 215–229. <https://doi.org/10.1007/s10705-020-10065-x>
- Martin, M., Dimassi, B., Roman Dobarco, M., Guenet, B., Arrouays, D., Angers, D.A., Soussana, J.F., Pellerin, S., 2021 (accepted). Feasibility of the 4 per 1000 aspirational target for soil carbon. A case study for France. *Global Change Biology*.
- Mary, B., Justes, E., 2012. Effet des cultures intermédiaires sur la séquestration du carbone et le bilan gaz à effet de serre. Justes, E. et al., Réduire les fuites de nitrate au moyen de cultures intermédiaires. <http://prodinra.inra.fr/record/168982>.



- Pellerin, S., Bamière, L., Angers, D., Béline, F., Benoît, M., Butault, J.P., 2013. How can French agriculture contribute to reducing greenhouse gas emissions? (Synopsis of the study report, INRA (France)).
- Pellerin, S., Bamière, L., Angers, D., Béline, F., Benoit, M., Butault, J.-P., Chenu, C., Colnenne-David, C., De Cara, S., Delame, N., Doreau, M., Dupraz, P., Faverdin, P., Garcia-Launay, F., Hassouna, M., Hénault, C., Jeuffroy, M.-H., Klumpp, K., Metay, A., Moran, D., Recous, S., Samson, E., Savini, I., Pardon, L., Chemineau, P., 2017. Identifying cost-competitive greenhouse gas mitigation potential of French agriculture. *Environmental Science & Policy* 77, 130–139. <https://doi.org/10.1016/j.envsci.2017.08.003>
- Pellerin, S., Bamière, L., Launay, C., Martin, R., Schiavo, M., Angers, D., Augusto, L., Balesdent, J., Doelsch, I.B., Bellassen, V., 2019. Stocker du carbone dans les sols français, quel potentiel au regard de l'objectif 4 pour 1000 et à quel coût? (Synthèse du rapport d'étude, INRA (France)).
- Peltre, C., Christensen, B.T., Dragon, S., Icard, C., Kätterer, T., Houot, S., 2012. RothC simulation of carbon accumulation in soil after repeated application of widely different organic amendments. *Soil Biology and Biochemistry* 52, 49–60. <https://doi.org/10.1016/j.soilbio.2012.03.023>

Germany

- Billen, N., Röder, C., Gaiser, T., Stahr, K., 2009. Carbon sequestration in soils of SW-Germany as affected by agricultural management—Calibration of the EPIC model for regional simulations. *Ecological Modelling* 220, 71–80. <https://doi.org/10.1016/j.ecolmodel.2008.08.015>
- Gaiser, T., Stahr, K., Billen, N., Mohammad, M.A.-R., 2008. Modeling carbon sequestration under zero tillage at the regional scale. I. The effect of soil erosion. *Ecological Modelling* 218, 110–120.
- Kanzler, M., Böhm, C., Freese, D., 2020. The development of soil organic carbon under young black locust (*Robinia pseudoacacia* L.) trees at a post-mining landscape in eastern Germany. *New Forests*. <https://doi.org/10.1007/s11056-020-09779-1>
- Poeplau, C., Zopf, D., Greiner, B., Geerts, R., Korvaar, H., Thumm, U., Don, A., Heidkamp, A., Flessa, H., 2018. Why does mineral fertilization increase soil carbon stocks in temperate grasslands? *Agriculture, ecosystems & environment* 265, 144–155.
- Wiesmeier, M., Hübner, R., Spörlein, P., Geuß, U., Hangen, E., Reischl, A., Schilling, B., Lützw, M. von, Kögel-Knabner, I., 2014. Carbon sequestration potential of soils in southeast Germany derived from stable soil organic carbon saturation. *Global Change Biology* 20, 653–665. <https://doi.org/10.1111/gcb.12384>
- Wiesmeier, M., Mayer, S., Burmeister, J., Hübner, R., Kögel-Knabner, I., 2020. Feasibility of the 4 per 1000 initiative in Bavaria: A reality check of agricultural soil management and carbon sequestration scenarios. *Geoderma* 369, 114333. <https://doi.org/10.1016/j.geoderma.2020.114333>

Hungary

- Nagy, Z., Pintér, K., Czóbel, Sz., Balogh, J., Horváth, L., Fóti, Sz., Barcza, Z., Weidinger, T., Csintalan, Zs., Dinh, N.Q., Grosz, B., Tuba, Z., 2007. The carbon budget of semi-arid grassland in a wet and a dry year in Hungary. *Agriculture, Ecosystems & Environment, The Greenhouse Gas Balance of Grasslands in Europe* 121, 21–29. <https://doi.org/10.1016/j.agee.2006.12.003>
- Pintér, K., Barcza, Z., Balogh, J., Czóbel, Sz., Csintalan, Zs., Tuba, Z., Nagy, Z., 2008. Interannual variability of grasslands' carbon balance depends on soil type. *COMMUNITY ECOLOGY* 9, 43–48. <https://doi.org/10.1556/ComEc.9.2008.S.7>



Ireland

- Abdalla, M., Hastings, A., Helmy, M., Prescher, A., Osborne, B., Lanigan, G., Forristal, D., Killi, D., Maratha, P., Williams, M., Rueangritsarakul, K., Smith, P., Nolan, P., Jones, M.B., 2014. Assessing the combined use of reduced tillage and cover crops for mitigating greenhouse gas emissions from arable ecosystem. *Geoderma* 223–225, 9–20. <https://doi.org/10.1016/j.geoderma.2014.01.030>
- Lanigan G., Forristal, D., Prescher, A.-K., 2012: Assessing the greenhouse gas budget of tillage mitigation options for arable systems. *Technology updates, Crops environment and Land Use. Agriculture and Food development authority, Teagasc.* https://www.teagasc.ie/media/website/publications/2012/GHG-from-ecotillage_5802.pdf
- Lanigan, G., Donnellan, T., Hanrahan, K., Carsten, P., Shalloo, L., Krol, D., Forristal, P.J., Farrelly, N., O'Brien, D., Ryan, M., Murphy, P., Caslin, B., Spink, J., Finnan, J., Boland, A., Upton, J., Richards, K.G., 2018. *An Analysis of Abatement Potential of Greenhouse Gas Emissions in Irish Agriculture 2021-2030 (Technical Report).* Teagasc.
- Simo, I., Schulte, R., O'Sullivan, L., Creamer, R., 2019. Digging deeper: Understanding the contribution of subsoil carbon for climate mitigation, a case study of Ireland. *Environmental Science & Policy* 98, 61–69. <https://doi.org/10.1016/j.envsci.2019.05.004>
- Torres-Sallan, G., Schulte, R., Lanigan, G.J., Byrne, K.A., Reidy, B., Simo, I., Six, J., Creamer, R.E., 2017a. Clay illuviation provides a longterm sink for C sequestration in subsoils. <https://doi.org/10.1038/srep45635>
- van Groenigen, K.J., Hastings, A., Forristal, D., Roth, B., Jones, M., Smith, P., 2011. Soil C storage as affected by tillage and straw management: An assessment using field measurements and model predictions. *Agriculture, Ecosystems & Environment* 140, 218–225. <https://doi.org/10.1016/j.agee.2010.12.008>

Italy

- Baldi, E., Cavani, L., Margon, A., Quartieri, M., Sorrenti, G., Marzadori, C., Toselli, M., 2018. Effect of compost application on the dynamics of carbon in a nectarine orchard ecosystem. *Science of The Total Environment* 637–638, 918–925. <https://doi.org/10.1016/j.scitotenv.2018.05.093>
- Barbera, V., Poma, I., Gristina, L., Novara, A., Egli, M., 2012. Long-term cropping systems and tillage management effects on soil organic carbon stock and steady state level of C sequestration rates in a semiarid environment. *Land Degrad. Dev.* 23, 82–91. <https://doi.org/10.1002/ldr.1055>
- Bleuler, M., Farina, R., Francaviglia, R., di Bene, C., Napoli, R., Marchetti, A., 2017. Modelling the impacts of different carbon sources on the soil organic carbon stock and CO₂ emissions in the Foggia province (Southern Italy). *Agricultural Systems* 157, 258–268. <https://doi.org/10.1016/j.agsy.2017.07.017>
- Bonciarelli, U., Onofri, A., Benincasa, P., Farneselli, M., Guiducci, M., Pannacci, E., Tosti, G., Tei, F., 2016. Long-term evaluation of productivity, stability and sustainability for cropping systems in Mediterranean rainfed conditions. *European Journal of Agronomy* 77, 146–155. <https://doi.org/10.1016/j.eja.2016.02.006>
- Borrelli, L., Colecchia, S., Troccoli, A., Caradonna, S., Papini, R., Ventrella, D., Dell'Abate, M.T., Galeffi, C., Tomasoni, C., Farina, R., 2011. Effectiveness of the GAEC standard of cross compliance Crop rotations in maintaining organic matter levels in soils. *Ital J Agronomy* 6, 8. <https://doi.org/10.4081/ija.2011.6.s1.e8>
- Cillis, D., Maestrini, B., Pezzuolo, A., Marinello, F., Sartori, L., 2018. Modeling soil organic carbon and carbon dioxide emissions in different tillage systems supported by precision agriculture



- technologies under current climatic conditions. *Soil and Tillage Research* 183, 51–59.
<https://doi.org/10.1016/j.still.2018.06.001>
- D’Avino, L., Di Bene, C., Farina, R., Razza, F., 2020. Introduction of Cardoon (*Cynara cardunculus* L.) in a Rainfed Rotation to Improve Soil Organic Carbon Stock in Marginal Lands. *Agronomy* 10, 946.
<https://doi.org/10.3390/agronomy10070946>
- Di Bene, C., Marchetti, A., Francaviglia, R., Farina, R., 2016. Soil organic carbon dynamics in typical durum wheat-based crop rotation of Southern Italy. *Ital J Agronomy* 11.
<https://doi.org/10.4081/ija.2016.763>
- Farina, R., Testani, E., Campanelli, G., Leteo, F., Napoli, R., Canali, S., Tittarelli, F., 2018. Potential carbon sequestration in a Mediterranean organic vegetable cropping system. A model approach for evaluating the effects of compost and Agro-ecological Service Crops (ASCs). *Agricultural Systems* 162, 239–248. <https://doi.org/10.1016/j.agsy.2018.02.002>
- Fiore, A., Lardo, E., Montanaro, G., Laterza, D., Loiudice, C., Berloco, T., Dichio, B., Xiloyannis, C., 2018. Mitigation of global warming impact of fresh fruit production through climate smart management. *Journal of Cleaner Production* 172, 3634–3643.
<https://doi.org/10.1016/j.jclepro.2017.08.062>
- Francaviglia, R., Coleman, K., Whitmore, A.P., Doro, L., Urracci, G., Rubino, M., Ledda, L., 2012. Changes in soil organic carbon and climate change – Application of the RothC model in agro-silvo-pastoral Mediterranean systems. *Agricultural Systems* 112, 48–54.
<https://doi.org/10.1016/j.agsy.2012.07.001>
- Iocola, I., Bassu, S., Farina, R., Antichi, D., Basso, B., Bindi, M., Dalla Marta, A., Danuso, F., Doro, L., Ferrise, R., Giglio, L., Ginaldi, F., Mazzoncini, M., Mula, L., Orsini, R., Corti, G., Pasqui, M., Seddaiu, G., Tomozeiu, R., Ventrella, D., Villani, G., Roggero, P.P., 2017. Can conservation tillage mitigate climate change impacts in Mediterranean cereal systems? A soil organic carbon assessment using long term experiments. *European Journal of Agronomy* 90, 96–107.
<https://doi.org/10.1016/j.eja.2017.07.011>
- Lugato, E., Berti, A., Giardini, L., 2006. Soil organic carbon (SOC) dynamics with and without residue incorporation in relation to different nitrogen fertilisation rates. *Geoderma* 135, 315–321.
<https://doi.org/10.1016/j.geoderma.2006.01.012>
- Mazzoncini, M., Antichi, D., Di Bene, C., Risaliti, R., Petri, M., Bonari, E., 2016. Soil carbon and nitrogen changes after 28 years of no-tillage management under Mediterranean conditions. *European Journal of Agronomy* 77, 156–165. <https://doi.org/10.1016/j.eja.2016.02.011>
- Mazzoncini, M., Sapkota, T.B., Barberi, P., Antichi, D., Risaliti, R., 2011. Long-term effect of tillage, nitrogen fertilization and cover crops on soil organic carbon and total nitrogen content. *Soil and Tillage Research* 114, 165–174. <https://doi.org/10.1016/j.still.2011.05.001>
- Mohamad, R.S., Verrastro, V., Al Bitar, L., Roma, R., Moretti, M., Al Chami, Z., 2016. Effect of different agricultural practices on carbon emission and carbon stock in organic and conventional olive systems. *Soil Res.* 54, 173. <https://doi.org/10.1071/SR14343>
- Mondini, C., Coleman, K., Whitmore, A.P., 2012. Spatially explicit modelling of changes in soil organic C in agricultural soils in Italy, 2001–2100: Potential for compost amendment. *Agriculture, Ecosystems & Environment* 153, 24–32. <https://doi.org/10.1016/j.agee.2012.02.020>
- Montanaro, G., Tuzio, A.C., Xylogiannis, E., Kolimenakis, A., Dichio, B., 2017. Carbon budget in a Mediterranean peach orchard under different management practices. *Agriculture, Ecosystems & Environment* 238, 104–113. <https://doi.org/10.1016/j.agee.2016.05.031>
- Novara, A., Minacapilli, M., Santoro, A., Rodrigo-Comino, J., Carrubba, A., Sarno, M., Venezia, G., Gristina, L., 2019. Real cover crops contribution to soil organic carbon sequestration in sloping vineyard. *Science of The Total Environment* 652, 300–306.
<https://doi.org/10.1016/j.scitotenv.2018.10.247>
- Triberti, L., Nastri, A., Giordani, G., Comellini, F., Baldoni, G., Toderi, G., 2008. Can mineral and organic fertilization help sequester carbon dioxide in cropland? *European Journal of Agronomy* 29,



13–20. <https://doi.org/10.1016/j.eja.2008.01.009>

Valkama, E., Kunyupiyeva, G., Zhapayev, R., Karabayev, M., Zhusupbekov, E., Perego, A., Schillaci, C., Sacco, D., Moretti, B., Grignani, C., Acutis, M., 2020. Can conservation agriculture increase soil carbon sequestration? A modelling approach. *Geoderma* 369, 114298. <https://doi.org/10.1016/j.geoderma.2020.114298>

Latvia

No studies available

Lithuania

Feiziene, D., Feiza, V., Karklins, A., Versulienė, A., Janusauskaite, D., Antanaitis, S., 2018. After-effects of long-term tillage and residue management on topsoil state in Boreal conditions. *European Journal of Agronomy* 94, 12–24.

Feiziene, D., Feiza, V., Slepėtienė, A., Liaudanskiene, I., Kadziene, G., Deveikyte, I., Vaideliene, A., 2011. Long-term influence of tillage and fertilization on net carbon dioxide exchange rate on two soils with different textures. *Journal of Environmental Quality* 40, 1787–1796.

Feizienė, D., Kadžienė, G., 2008. The influence of soil organic carbon, moisture and temperature on soil surface CO₂ emission in the 10th year of different tillage-fertilisation management. *Zemdirbyste-Agriculture* 95, 29–45.

Jokubauskaite, I., Karčauskienė, D., Slepėtienė, A., Repsiene, R., Amaleviciute, K., 2016. Effect of different fertilization modes on soil organic carbon sequestration in acid soils. *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science* 66, 647–652.

Skinulienė, L., Bogužas, V., Aleinikovienė, J., Steponavičienė, V., Sinkevičienė, A., 2018. Effect of 50-year crop rotations on soil organic carbon content.

Lithuania's Greenhouse Gas Inventory Report 2020. [online][15.04.2020]. Available at: http://klimatas.gamta.lt/files/NIR_2020%2004%2015.pdf

Netherlands

Akker, J. J., Hoving, I., Hendriks, R., & Knotters, M. (2018). *Onderwaterdrains zijn effectief* (No. 2922). Wageningen Environmental Research.

Koopmans, C., Timmermans, B., de Haan, J., van Opheusden, M., Selin Noren, I., Slier, T., & Wagenaar, J. P. (2020). *Evaluatie van maatregelen voor het vastleggen van koolstof in minerale gronden 2019-2023. Voortgangsrapportage april 2020*. Louis Bolk Instituut.

Lesschen, J. P., Heesmans, H. I. M., Mol-Dijkstra, J. P., van Doorn, A. M., Verkaik, E., van den Wyngaert, I. J. J., & Kuikman, P. J. (2012). *Mogelijkheden voor koolstofvastlegging in de Nederlandse landbouw en natuur* (No. 2396). Alterra.

Koopmans, C., Timmermans, B., Wagenaar, J.P., van 't Hull, J., Hanegraaf, M., & de Haan, J., (2019). *Evaluatie van maatregelen voor het vastleggen van koolstof. Resultaten uit Lange Termijn Experimenten (LTE's)*. Louis Bolk Instituut.

Van den Born, G. J., Kragt, F., Henkens, D., Rijken, B., Van Bommel, B., Van der Sluis, S., ... & van den Akker, J. (2016). *Dalende bodems, stijgende kosten: mogelijke maatregelen tegen veenbodemdaling in het landelijk en stedelijk gebied: beleidsstudie* (No. 1064). Planbureau voor de Leefomgeving.

Norway

Bøe, F., Bechmann, M., Øgaard, A.F., Sturite, I., Brandsæter, L.O., 2019. *Fangvekstenes økosystemtjenester – Kunnskapsstatus om effekter av fangvekster* (Nibio No. 5). Norway.

Cuvaradic, M., Tveitnes, S., Krogstad, T., Lombnæs, P., 2004. Long-term effects of crop rotation and



different fertilization systems on soil fertility and productivity. *Acta Agriculturae Scandinavica, Section B-Soil & Plant Science* 54, 193–201.

- Grønlund, A., de Zarruk, K.K., Rasse, D., Riley, H., Klakegg, O., Nystuen, I., 2008. Kunnskapsstatus for utslipp og binding av karbon i jordbruksjord. *Bioforsk rapport* 3, 47.
- Rasse, D., Økland, I.H., Bárcena, T.G., Riley, H., Martinsen, V., Sturite, I., Joner, E., O’Toole, A., Øpstad, S., Cottis, T., 2019. Muligheter og utfordringer for økt karbonbinding i jordbruksjord. *NIBIO rapport*.
- Singh, B.R., Lal, R., 2005. The potential of soil carbon sequestration through improved management practices in Norway. *Environment, Development and Sustainability* 7, 161–184.

Poland

- Faber A., Jarosz Z. (2018). Modeling of Soil Organic Carbon Balance and Greenhouse Gas Emissions on a Regional Scale and in Poland (in Polish). *Problems of World Agriculture*, 18(3), 102–112; DOI: 10.22630/PRS.2018.18.3.70
- Faber, A., Jarosz, Z., Łopata, A., Siebielec, G. (2012). Assessment of changes in organic carbon content in soils based on monitoring data on arable soil chemistry in Poland. *Studia i Raporty IUNG-PIB*, 46(20), 9-20. (in Polish)
- Krasuska, E., Faber, A., Jarosz, Z., Kaczyński, R. (2016). Modelling soil organic carbon sequestration under crops for biofuels in Poland. *Polish Journal of Agronomy*, 27, 126-136.
- Borzecka-Walker M., Faber A. & Borek R. (2008). Evaluation of carbon sequestration in energetic crops (*Miscanthus* and coppice willow). *Int. Agrophysics*, No 22, 185-190.
- Borzęcka-Walker M, Faber A, Mizak K, Pudełko R, Syp A (2011) Soil carbon sequestration under bioenergy crops in Poland. In: *Principles, Application and Assessment in Soil Science* (ed. EB Özkarova Güngör), pp. 151– 166. InTech, Rijeka, Croatia.

Portugal

- Borges, O., Raimundo, F., Coutinho, J., Gonçalves, B., Oliveira, I., Martins, A., Madeira, M., 2018. Carbon fractions as indicators of organic matter dynamics in chestnut orchards under different soil management practices. *Agroforestry systems* 92, 301–310.
- Carvalho, M., Barsch, G., Calado, J.M.G., Barros, J.F.C., 2012. Long term effect of tillage system and crop residue management on soil carbon content of a Luvisol under rainfed Mediterranean conditions. *Agrociencia-Sitio en Reparación* 16 3, 183–187.
- Pereira Costa, T., Pina, A., Canaveira, P., Amaro, A., Borges, M., 2020. Portuguese National Inventory Report on Greenhouse Gases, 1990 – 2018. Portuguese Environment Agency. Annex E., Amadora.
- Sil, Â., Fonseca, F., Gonçalves, J., Honrado, J., Marta-Pedroso, C., Alonso, J., Ramos, M., Azevedo, J.C., 2017. Analysing carbon sequestration and storage dynamics in a changing mountain landscape in Portugal: insights for management and planning. *International Journal of Biodiversity Science, Ecosystem Services & Management* 13, 82–104. <https://doi.org/10.1080/21513732.2017.1297331>
- Teixeira, R., Domingos, T., Valada, T., Martins, H., Calouro, F., 2020. Methodological note concerning the calculation of carbon sequestration in areas with sown biodiverse pastures.
- Teixeira, R.F.M., Domingos, T., Costa, A., Oliveira, R., Farropas, L., Calouro, F., Barradas, A.M., Carneiro, J., 2011. Soil organic matter dynamics in Portuguese natural and sown rainfed grasslands. *Ecological Modelling* 222, 993–1001.
- Valada, T., Castro Ribeiro, M., Martins, H., Domingos, T., 2020. Methodological note concerning the calculation of carbon sequestration in areas where harrowing was replaced by less disruptive methods for shrub control. In *PEA (2020) Portuguese National Inventory Report on Greenhouse Gases, 1990 – 2018*. Amadora: Portuguese Environment Agency. Annex F.



Valada, T., Teixeira, R., Martins, H., Ribeiro, M., Domingos, T., 2012. Grassland management options under Kyoto Protocol Article 3.4. Sustainable Mediterranean Grasslands and Their Multi-Functions; Options Méditerranéennes; CIHEAM/FAO/ENMP/SPPF: Zaragoza, Spain 53–56.

Slovakia

Barančíková, G., Halas, J., Koco, Š., Gutteková, M., 2011b: Influence of soil protection system on development of organic carbon stock. *Soil Science Agrochemistry and Ecology*, vol. XLV, No.1-4, pp. 27-31.

Barančíková, G., Skalský, R., Koco, Š., Halas, J., Tarasovičová, Z., Nováková, M. 2014. Farm-level modelling of soil organic carbon sequestration under climate and land use change. Halldórsson, G., Bampa, F., Porsteinsdóttir, A.B. (eds.) *Soil carbon sequestration for climate food security and ecosystem services*. Luxembourg: Publications Office of the European Union, 2014, pp. 94-100.

Halas, J., Makovníková, J., Barančíková, G., Tarasovičová, Z., Koco, Š., Skalský, R., 2018: Hodnotenie zmien obsahu pôdneho organického uhlíka (POC) podľa pôdnych typov (Evaluation of soil organic carbon (SOC) changes according soil types) . In: Ilavská, B. (Ed.): *Zborník abstraktov – pedologické dni 2018 (Proceedings – Pedological days)*, Bratislava, 12.-14.9.2018, str.47, NPPC-VÚPOP Bratislava, Societas Pedologica Slovaca, 2018. ISBN 978-80-8163-027-9

Slovenia

Kaurin A., Mihelič R., Kastelec D., Schloter M., Suhadolc M., Grčman H. 2015. Consequences of minimum soil tillage on abiotic soil properties and composition of microbial communities in a shallow Cambisol originated from fluvioglacial deposits. *Biology and Fertility of Soils*, 51: 923-933

Vidrih M. 2006. Carbon sequestration in grazed sward of high karst. Doctoral dissertation. Ljubljana, Univerza v Ljubljani, Biotehniška fakulteta: 102 p.

Vončina A. 2015. Carbon balance of fire prone karst grasslands. Doctoral dissertation. Ljubljana, Univerza v Ljubljani, Biotehniška fakulteta: 86 p.

Spain

Álvaro-Fuentes, J., Plaza-Bonilla, D., Arrúe, J.L., Lampurlanés, J., Cantero-Martínez, C., 2014. Soil organic carbon storage in a no-tillage chronosequence under Mediterranean conditions. *Plant Soil* 376, 31–41. <https://doi.org/10.1007/s11104-012-1167-x>

Calleja-Cervantes, M.E., Aparicio-Tejo, P.M., Villadas, P.J., Irigoyen, I., Irañeta, J., Fernández-González, A.J., Fernández-López, M., Menéndez, S., 2017. Rational application of treated sewage sludge with urea increases GHG mitigation opportunities in Mediterranean soils. *Agriculture, Ecosystems & Environment*, Quantification and mitigation of greenhouse gas emissions in Mediterranean cropping systems 238, 114–127. <https://doi.org/10.1016/j.agee.2016.09.021>

Carbonell-Bojollo, R., González-Sánchez, E.J., Veróz-González, O., Ordóñez-Fernández, R., 2011. Soil management systems and short term CO₂ emissions in a clayey soil in southern Spain. *Science of The Total Environment* 409, 2929–2935. <https://doi.org/10.1016/j.scitotenv.2011.04.003>

Garcia-Pausas, J., Rabissi, A., Rovira, P., Romanyà, J., 2017. Organic Fertilisation Increases C and N Stocks and Reduces Soil Organic Matter Stability in Mediterranean Vegetable Gardens. *Land Degradation & Development* 28, 691–698. <https://doi.org/10.1002/ldr.2569>

González-Sánchez, E.J., Ordóñez-Fernández, R., Carbonell-Bojollo, R., Veroz-González, O., Gil-Ribes, J.A., 2012. Meta-analysis on atmospheric carbon capture in Spain through the use of conservation agriculture. *Soil and Tillage Research* 122, 52–60. <https://doi.org/10.1016/j.still.2012.03.001>

Lammerding, D.M., Hontoria, C., Tenorio, J.L., Walter, I., 2011. Mediterranean Dryland Farming: Effect of Tillage Practices on Selected Soil Properties. *Agronomy Journal* 103, 382–389. <https://doi.org/10.2134/agronj2010.0210>



- López-Bellido, L., López-Bellido, R., Fernández-García, P., Muñoz-Romero, V., Lopez-Bellido, F.J., 2020. Carbon storage in a rainfed Mediterranean vertisol: Effects of tillage and crop rotation in a long-term experiment. *European Journal of Soil Science* 71, 472–483. <https://doi.org/10.1111/ejss.12883>
- Moreno-García, M., Repullo-Ruibérriz de Torres, M.Á., González-Sánchez, E.J., Ordóñez-Fernández, R., Veroz-González, Ó., Carbonell-Bojollo, R.M., 2020. Methodology for estimating the impact of no tillage on the 4perMille initiative: The case of annual crops in Spain. *Geoderma* 371, 114381. <https://doi.org/10.1016/j.geoderma.2020.114381>
- Parras-Alcántara, L., Lozano-García, B., 2014. Conventional tillage versus organic farming in relation to soil organic carbon stock in olive groves in Mediterranean rangelands (southern Spain). *Solid Earth* 5, 299–311. <https://doi.org/10.5194/se-5-299-201>

Sweden

- Bolinder, M.A., Kätterer, T., Andrén, O., Ericson, L., Parent, L.-E., Kirchmann, H., 2010. Long-term soil organic carbon and nitrogen dynamics in forage-based crop rotations in Northern Sweden (63–64 N). *Agriculture, ecosystems & environment* 138, 335–342.
- Börjesson, G., Kätterer, T., 2018a. Soil fertility effects of repeated application of sewage sludge in two 30-year-old field experiments. *Nutrient cycling in agroecosystems* 112, 369–385.
- Börjesson, G., Kätterer, T., 2018b. Soil fertility effects of repeated application of sewage sludge in two 30-year-old field experiments. *Nutr Cycl Agroecosyst* 112, 369–385. <https://doi.org/10.1007/s10705-018-9952-4>
- Eckersten, H., Bolinder, M., Johnsson, H., Kätterer, T., Mårtensson, K., Collentine, D., Tidåker, P., Marstorp, H., 2017. Nitrogen leaching and soil organic carbon sequestration of a Barley crop with improved N use efficiency—A regional case study. *Acta Agriculturae Scandinavica, Section B—Soil & Plant Science* 67, 615–627.
- Haddaway, N.R., Hedlund, K., Jackson, L.E., Kätterer, T., Lugato, E., Thomsen, I.K., Jørgensen, H.B., Söderström, B., 2015. What are the effects of agricultural management on soil organic carbon in boreo-temperate systems? *Environmental Evidence* 4, 23.
- Karlsson, A.-B., Boström, B., Jernbäcker, E., Lundblad, M., Mjureke, D., 2020. Vägen till en klimatpositiv framtid. *SOU 2020: Betänkande från Klimatpolitiska vägvalsutredningen (M 2018:07)*.
- Kätterer, T., Andersson, L., Andrén, O., Persson, J., 2008. Long-term impact of chronosequential land use change on soil carbon stocks on a Swedish farm. *Nutrient Cycling in Agroecosystems* 81, 145.
- Kätterer, T., Bolinder, M.A., Andrén, O., Kirchmann, H., Menichetti, L., 2011. Roots contribute more to refractory soil organic matter than above-ground crop residues, as revealed by a long-term field experiment. *Agriculture, Ecosystems & Environment* 141, 184–192.
- Kätterer, T., Börjesson, G., Kirchmann, H., 2014. Changes in organic carbon in topsoil and subsoil and microbial community composition caused by repeated additions of organic amendments and N fertilisation in a long-term field experiment in Sweden. *Agriculture, Ecosystems & Environment* 189, 110–118. <https://doi.org/10.1016/j.agee.2014.03.025>
- Menichetti, L., Ekblad, A., Kätterer, T., 2015. Contribution of roots and amendments to soil carbon accumulation within the soil profile in a long-term field experiment in Sweden. *Agriculture, Ecosystems & Environment* 200, 79–87.
- Meurer, K.H., Haddaway, N.R., Bolinder, M.A., Kätterer, T., 2018. Tillage intensity affects total SOC stocks in boreo-temperate regions only in the topsoil—A systematic review using an ESM approach. *Earth-Science Reviews* 177, 613–622.
- Poeplau, C., Aronsson, H., Myrbeck, Å., Kätterer, T., 2015a. Effect of perennial ryegrass cover crop on soil organic carbon stocks in southern Sweden. *Geoderma Regional* 4, 126–133.
- Poeplau, C., Bolinder, M., Eriksson, J., Lundblad, M., Kätterer, T., 2015b. Positive trends in organic carbon storage in Swedish agricultural soils due to unexpected socio-economic drivers.



- Biogeosciences 12: 3241–3251.
- Poeplau, C., Kätterer, T., Bolinder, M.A., Börjesson, G., Berti, A., Lugato, E., 2015c. Low stabilization of aboveground crop residue carbon in sandy soils of Swedish long-term experiments. *Geoderma* 237, 246–255.
- Vägen till en klimatpositiv framtid. SOU 2020: Betänkande från Klimatpolitiska vägvalsutredningen (M 2018:07)., 2020.
- Wikström, L., 2019. Så klarar det svenska jordbruket klimatmålen: en delrapport från IVA-projektet Vägval för klimatet.

Switzerland

- Ammann, C., Neftel, A., Joher, M., Fuhrer, J., Leifeld, J., 2020. Effect of management and weather variations on the greenhouse gas budget of two grasslands during a 10-year experiment. *Agriculture, Ecosystems & Environment* 292, 106814. <https://doi.org/10.1016/j.agee.2019.106814>
- Anken, T., Weisskopf, P., Zihlmann, U., Forrer, H., Jansa, J., Perhacova, K., 2004. Long-term tillage system effects under moist cool conditions in Switzerland. *Soil and Tillage Research, Soil Quality as an Indicator of Sustainable Tillage Practices* 78, 171–183. <https://doi.org/10.1016/j.still.2004.02.005>
- Beuttler, C., Keel, S.G., Leifeld, J., Schmid, M., Berta, N., Gutknecht, V., Wohlgemuth, N., Brodmann, U., Stadler, Z., Tinibaev, D., Wlodarczak, D., Honegger, M., Stettler, C., 2019. The Role of Atmospheric Carbon Dioxide Removal in Swiss Climate Policy – Fundamentals and Recommended Actions., Report by Risk Dialogue Foundation. Commissioned by the Federal Office for the Environment, Bern., Bern.
- Hermle, S., Anken, T., Leifeld, J., Weisskopf, P., 2008. The effect of the tillage system on soil organic carbon content under moist, cold-temperate conditions. *Soil and Tillage Research* 98, 94–105. <https://doi.org/10.1016/j.still.2007.10.010>
- Keel, S.G., Anken, T., Büchi, L., Chervet, A., Fliessbach, A., Flisch, R., Huguenin-Elie, O., Mäder, P., Mayer, J., Sinaj, S., Sturny, W., Wüst-Galley, C., Zihlmann, U., Leifeld, J., 2019. Loss of soil organic carbon in Swiss long-term agricultural experiments over a wide range of management practices. *Agriculture, Ecosystems & Environment* 286, 106654. <https://doi.org/10.1016/j.agee.2019.106654>
- Krauss, M., Ruser, R., Müller, T., Hansen, S., Mäder, P., Gattinger, A., 2017. Impact of reduced tillage on greenhouse gas emissions and soil carbon stocks in an organic grass-clover ley - winter wheat cropping sequence. *Agriculture, Ecosystems & Environment* 239, 324–333. <https://doi.org/10.1016/j.agee.2017.01.029>
- Leifeld, J., Bassin, S., Fuhrer, J., 2003. Carbon stocks and carbon sequestration potentials in agricultural soils in Switzerland. *Agroscope, FAL-Schriftenreihe* 44, Zürich. - Google Search (Schriftreihe FAL No. 44). Eidgenössische Forschungsanstalt für Agrarökologie und Landbau, FAL Reckenholz, Zürich.
- Leifeld, J., Bassin, S., Fuhrer, J., 2005. Carbon stocks in Swiss agricultural soils predicted by land-use, soil characteristics, and altitude. *Agriculture, Ecosystems & Environment* 105, 255–266. <https://doi.org/10.1016/j.agee.2004.03.006>
- Martínez, I., Chervet, A., Weisskopf, P., Sturny, W.G., Etana, A., Stettler, M., Forkman, J., Keller, T., 2016. Two decades of no-till in the Oberacker long-term field experiment: Part I. Crop yield, soil organic carbon and nutrient distribution in the soil profile. *Soil and Tillage Research* 163, 141–151. <https://doi.org/10.1016/j.still.2016.05.021>
- Oberholzer, H.R., Leifeld, J., Mayer, J., 2014. Changes in soil carbon and crop yield over 60 years in the Zurich Organic Fertilization Experiment, following land-use change from grassland to cropland. *Journal of Plant Nutrition and Soil Science* 177, 696–704.



<https://doi.org/10.1002/jpln.201300385>

Turkey

- Akça, M.O., Namli, A., 2015. Effects of poultry litter biochar on soil enzyme activities and tomato, pepper and lettuce plants growth. *Eurasian Journal of Soil Science* 4, 161.
- Ortas, I., Akpınar, C., Lal, R., 2013. Long-Term Impacts of Organic and Inorganic Fertilizers on Carbon Sequestration in Aggregates of an Entisol in Mediterranean Turkey. *Soil Science* 178, 12–23. <https://doi.org/10.1097/SS.0b013e3182838017>

United Kingdom

- Bellamy, P. H., Loveland, P. J., Bradley, R. I., Lark, R. M., Kirk, G. J. D.: Carbon losses from all soils across England and Wales 1978–2003. *Nature* 437, 245–248, 2005.
- Bradley, R. I., R. Milne, et al. (2005). A soil carbon and land-use database for the United Kingdom. *Soil Use and Management* 21(004): 363-369.
- Brown, P., Cardenas, L., Choudrie, S., Jones, L., Karagianni, E., MacCarthy, J., Passant, N., Richmond, B., Smith, H., 2020. UK Greenhouse Gas Inventory, 1990 to 2018: Annual Report for submission under the Framework Convention on Climate Change - Defra, UK. Department for Environment, Food and Rural Affairs (Defra), Nobel House, 17 Smith Square, London SW1P 3JR helpline@defra.gsi.gov.uk.
- Dawson, J.J.C., Smith, P., 2007. Carbon losses from soil and its consequences for land-use management. *Sci.Total Env.* 382, 165–190. <https://doi.org/10.1016/j.scitotenv.2007.03.023>
- Falloon, P., Powlson, D., Smith, P., 2004. Managing field margins for biodiversity and carbon sequestration: a Great Britain case study. *Soil Use and Management* 20, 240–247. <https://doi.org/10.1111/j.1475-2743.2004.tb00364.x>
- Fornara, D., Olave, R., Higgins, A., 2020. Evidence of low response of soil carbon stocks to grassland intensification. *Agriculture, Ecosystems & Environment* 287, 106705.
- Fornara, D.A., Banin, L., Crawley, M.J., 2013. Multi-nutrient vs. nitrogen-only effects on carbon sequestration in grassland soils. *Global Change Biology* 19, 3848–3857.
- Fornara, D.A., Steinbeiss, S., McNamara, N.P., Gleixner, G., Oakley, S., Poulton, P.R., Macdonald, A.J., Bardgett, R.D., 2011. Increases in soil organic carbon sequestration can reduce the global warming potential of long-term liming to permanent grassland. *Global Change Biology* 17, 1925–1934.
- Fornara, D.A., Wasson, E.-A., Christie, P., Watson, C.J., 2016. Long-term nutrient fertilization and the carbon balance of permanent grassland: any evidence for sustainable intensification? *Biogeosciences* 13, 4975.
- Poulton, P., Johnston, J., Macdonald, A., White, R., Powlson, D., 2018. Major limitations to achieving “4 per 1000” increases in soil organic carbon stock in temperate regions: Evidence from long-term experiments at Rothamsted Research, United Kingdom. *Global change biology* 24, 2563–2584.
- Powlson, D.S., Bhogal, A., Chambers, B.J., Coleman, K., Macdonald, A.J., Goulding, K.W.T., Whitmore, A.P., 2012. The potential to increase soil carbon stocks through reduced tillage or organic material additions in England and Wales: a case study. *Agriculture, Ecosystems & Environment* 146, 23–33
- Smith P, Martino D, Cai Z et al. (2008) Greenhouse gas mitigation in agriculture. *Philosophical Transactions of the Royal Society B: Biological Sciences*, 363, 789–813
- Smith, P. 2014. Do grasslands act as a perpetual sink for carbon? *Global Change Biol.* 20, 2708–2711



B List of references organic soils

Austria

No studies available

Belgium

No studies available

Czech Republic

No studies available

Denmark

No studies available were mitigation measures are described.

Petersen, S.O., Hoffmann, C.C., Schäfer, C.-M., Blicher-Mathiesen, G., Elsgaard, L., Kristensen, K., Larsen, S.E., Torp, S.B., Greve, M.H., 2012. Annual emissions of CH₄ and N₂O, and ecosystem respiration, from eight organic soils in Western Denmark managed by agriculture. Biogeosciences 9, 403.

Taghizadeh-Toosi, A., Elsgaard, L., Clough, T.J., Labouriau, R., Ernstsén, V., Petersen, S.O., 2019. Regulation of N₂O emissions from acid organic soil drained for agriculture. Biogeosciences 16, 4555–4575. <https://doi.org/10.5194/bg-16-4555-2019>

Estonia

No studies available

Finland

No studies available

France

No studies available

Germany

Tiemeyer, B., Freibauer, A., Borraz, E. A., Augustin, J., Bechtold, M., Beetz, S., . . . Fiedler, S. (2020). A new methodology for organic soils in national greenhouse gas inventories: Data synthesis, derivation and application. *J Ecological Indicators*, 109, 105838.

Tiemeyer et al. 2016. High emissions of greenhouse gases from grasslands on peat and other organic soils. *Global Change Biology* (2016) 22, 4134–4149, doi: 10.1111/gcb.13303

Hungary

No studies available

Ireland

Paul, C., Fealy, R., Fenton, O., Lanigan, G., O'Sullivan, L., Schulte, R.P.O., 2018. Assessing the role of artificially drained agricultural land for climate change mitigation in Ireland. *Environmental Science & Policy* 80, 95–104. <https://doi.org/10.1016/j.envsci.2017.11.004>



- Renou-Wilson, F., Barry, C., Müller, C., Wilson, D., 2014. The impacts of drainage, nutrient status and management practice on the full carbon balance of grasslands on organic soils in a maritime temperate zone. *Biogeosciences Discussions* 11, 5557–5612. <https://doi.org/10.5194/bgd-11-5557-2014>
- Wilson, D., Blain, D., Couwenberg, J., Evans, C.D., Murdiyarso, D., Page, S.E., Renou-Wilson, F., Rieley, J.O., Sirin, A., Strack, M., 2016. Greenhouse gas emission factors associated with rewetting of organic soils. *Mires and Peat* 17.
- Wilson, D., Müller, C., Renou-Wilson, F., 2014. Carbon emissions and removals from Irish peatlands: present trends and future mitigation measures. *Irish Geography* 46, 1–23. <https://doi.org/10.2014/igj.v46i1.261>
- Wilson, D., Renou-Wilson, F., Farrell, C., Bullock, C., Müller, C., 2012. Carbon restore—the potential of restored Irish peatlands for carbon uptake and storage. *The Potential of Peatlands for Carbon Sequestration (2007-CCRP-1.6)*. CCRP Report, Prepared for the Environmental Protection Agency. URL: www.epa.ie/downloads/pubs/research/climate/CCRP_15_web.pdf.

Italy (studies considering GHG emissions and mitigation; however, soils mentioned are not organic soils by definition)

- Bertora, C., Cucu, M.A., Lerda, C., Peyron, M., Bardi, L., Gorra, R., Sacco, D., Celi, L., Said-Pullicina, D., 2018. Dissolved organic carbon cycling, methane emissions and related microbial populations in temperate rice paddies with contrasting straw and water management. *Agric. Ecosyst. Environ. Appl. Soil Ecol.* 265, 1292–1306. <https://doi.org/10.1016/j.agee.2018.06.004>
- ISPRA, 2020. Italian Greenhouse Gas Inventory 1990-2018. National Inventory Report 2020. Available online at <https://www.isprambiente.gov.it/it/pubblicazioni/rapporti/italian-greenhouse-gas-inventory-1990-2018.-national-inventory-report-2020>
- Lagomarsino, A., Agnelli, A.E., Linqvist, B.A., Adviento-Borbe, M.A.A., Agnelli, A., Gavina, G., Ravaglia, S., Ferrara, R.M., 2016. Alternate wetting and drying of rice reduced CH₄ emissions but triggered N₂O peaks in a clayey soil of central Italy. *Pedosphere* 26, 533–548. [https://doi.org/10.1016/S1002-0160\(15\)60063-7](https://doi.org/10.1016/S1002-0160(15)60063-7)
- Mazza, G., Agnelli, A.E., Orasen, G., Gennaro, M., Valè, G., Lagomarsino, A., 2016. Reduction of Global Warming Potential from rice under alternate wetting and drying practice in a sandy soil of northern Italy. *Italian Journal of Agrometeorology-Rivista Italiana di Agrometeorologia* 21(2), 35-44. <https://doi.org/10.19199/2016.2.2038-5625.035>
- Meijide, A., Gruening, C., Goded, I., Seufert, G., Cescatti, A., 2017. Water management reduces greenhouse gas emissions in a Mediterranean rice paddy field. *Agric. Ecosyst. Environ.* 238, 168–178. <https://doi.org/10.1016/j.agee.2016.08.017>
- Miniotti, E.F., Romani, M., Said-Pullicino, D., Facchi, A., Bertora, C., Peyron, M., Sacco, D., Bischetti, G.B., Lerda, C., Tenni, D., Gandolfi, C., Celi, L., 2016. Agroenvironmental sustainability of different water management practices in temperate rice agro-ecosystems. *Agric. Ecosyst. Environ.* <https://doi.org/10.1016/j.agee.2016.02.010>.
- Peyron, M., Bertora, C., Pelissetti, S., Said-Pullicino, D., Celi, L., Miniotti, E., Romani, M., Sacco, D., 2016. Greenhouse gas emissions as affected by different water management practices in temperate rice paddies. *Agric. Ecosyst. Environ.* 232, 17–28. <https://doi.org/10.1016/j.agee.2016.07.021>

Latvia

No contribution



Lithuania

No studies available

Netherlands

Van den Akker, J. J. H., Massop, H. T. L., & Rietra, R. P. J. J. (2018). Potentiële emissiereductie broeikasgassen Fries veenweidegebied: waterhuishoudkundige en bodembeheermaatregelen om broeikasemissies te beperken (No. 2905). Wageningen Environmental Research

Van den Born, G. J., Kragt, F., Henkens, D., Rijken, B., Van Bommel, B., Van der Sluis, S., ... & van den Akker, J. (2016). Dalende bodems, stijgende kosten: mogelijke maatregelen tegen veenbodemdaling in het landelijk en stedelijk gebied: beleidsstudie (No. 1064). Planbureau voor de Leefomgeving.

Norway

No studies available

Poland

No studies available

Portugal

No studies available

Slovakia

No studies available

Slovenia

Pal L. 2011. The potential of Ljubljana marsh soil for mineral nitrogen removal. Dissertation thesis. Ljubljana, University of Ljubljana, Biotechnical faculty: 84 p., <http://www.ljubljanskobarje.si/uploads/datoteke/PAL%20LEVIN%20DR9.pdf>

Spain

No studies available

Sweden

Berglund, Ö. and Berglund, K. (2010) Distribution and cultivation intensity of agricultural peat and gyttja soils in Sweden and estimation of greenhouse gas emissions from cultivated peat soils. Geoderma Volume 154, Issues 3–4, 15 January 2010, Pages 173-180. <https://doi.org/10.1016/j.geoderma.2008.11.035>

Örjan Berglund, Kerstin Berglund, Sabine Jordan, Lisbet Norberg (2019) Carbon capture efficiency, yield, nutrient uptake and trafficability of different grass species on a cultivated peat soil, CATENA, Volume 173, 2019, Pages 175-182, ISSN 0341-8162,

Switzerland

Leifeld, J., Bassin, S., Fuhrer, J., 2003. Carbon stocks and carbon sequestration potentials in agricultural soils in Switzerland. Agroscope, FAL-Schriftenreihe 44, Zürich. - Google Search (Schriftreihe FAL No. 44). Eidgenössische Forschungsanstalt für Agrarökologie und Landbau, FAL Reckenholz, Zürich.

Turkey

No studies available



United Kingdom

Sozanska-Stanton, M., Carey, P.D., Griffiths, G.H., Vogiatzakis, I.N., Treweek, J., Butcher, B., Charlton, M.B., Keenleyside, C., Arnell, N.W., Tucker, G. and Smith, P., 2016. Balancing conservation and climate change—a methodology using existing data demonstrated for twelve UK priority habitats. *Journal for nature conservation*, 30, pp.76-89.

Evans, C., 2017. Implementation of an emission inventory for UK peatlands. Report to the Department for Business, Energy and Industrial Strategy, Centre for Ecology and Hydrology, Bangor. 88pp.



C List of finished and ongoing projects

Austria:

1. CASAS: CARbon Sequestration in Austrian Soils ONGOING (1.9.2019-31.8.2022). Document: Austrian Agency for Health and Food Safety (2019): Form Part A 3 - Proposal. 11th ACRP Call.
2. ACRP (Austrian Climate Research Programme) project PeatGov-Austria "Governance options for climate smart agriculture on Austrian peatlands - Specific support for Austria's policymakers" (no publication/report available yet).
3. CRP project "PeatGov-Austria" Austrian peat soils under agriculture are going to be mapped. A best estimate (no measurements) is the goal. In the APCC report quantitative estimates for reducing GHG emissions will be published.

Belgium

4. Investigation on two nature-based negative emission technologies: enhance weathering and biochar. Prof. E. Verbruggen; University of Antwerp.
5. BASTA-project; dr. Bart Vandecasteele; ILVO (Flanders, Belgium): Research on the greenhouse gas (GHG) emission mitigation potential of biochar during compost production and the GHG mitigation potential of this mixture after incorporation in soil.

Czech Republic

6. Research project NAZV QJ1320122 Optimization of the afforestation management of the agricultural lands in relation to enhancement of the landscape retention potential. (2013 - 2017).
7. Research project TACR TD03000087 Interactive evaluation of carbon sequestration in agricultural landscape (2016 - 2017), Evaluation of environmentally friendly practices within the RDP (rural development program) - impacts of interventions on the state of organic matter, biological activity in the soil and soil erodibility (2019).
8. Research project NAZV QK1910232 - The optimization of subsidy title for agricultural land afforestation (2019 - 2023),
9. Research project QK1910299 - Sustainable management of natural resources with emphasis on non-production and production ability of soil (2019 - 2023). General information about projects and its results are available on <https://starfos.tacr.cz/>

Estonia

Running project of Estonian Agricultural Research Centre:

10. The impact of no-till on soil properties
11. Peat and eroded soil in the soil protection measure (comparing arable fields and grasslands and changes in them during 6 years)
12. Fertility research to study SOC changes in the arable field depending on different types of subsidies (organic, environmentally friendly, single area payment)

Finland

13. Multi-benefit solutions to climate-smart agriculture, MULTA: <https://carbonaction.org/en-stn-multa/>

France

14. Stocker du carbone dans les sols français: <https://inra-dam-front-resources-cdn.wedia-group.com/ressources/afile/487878-58615-resource-etude-4-pour-1000-synthese-en-francais-pdf.pdf>
15. DEDYCAS: Soil carbon, a depth-dependent dynamic system: new concepts, measurements



- and modelling
16. StoreSoilC: Potential and sustainability of carbon storage in agricultural soils (2017-2021)
 17. NanoSoilC: Organo-mineral interactions: nanoscale mechanisms for carbon storage in soils (2017 - 2021)
 18. DSCATT: Agricultural Intensification and Dynamics of Soil Carbon Sequestration in Tropical and Temperate Farming Systems (2019-2023)
 19. CarSoilE: Construction of a methodology and a frame of reference on carbon fluxes in agricultural soils in cattle breeding territories (2018 – 2021)
 20. BRAWO: enhance fundamental knowledge of mechanisms involved in the degradation of woody litter. It will also contribute to the global effort for enhancing carbon sequestration in soil by assessing the potential of altered lignin for carbon sequestration and by developing an innovative microbial model of soil carbon dynamics.
 21. CLAND: Convergence Institute to perform in the next decade the research urgently needed on land-management solutions for managing the ecological and energy transitions of the 21st century (2017 – 2027)
 22. SUPRA: Develop new knowledge on carbon storage in urban soils in relation to climate change (2017 – 2020)
 23. PROTERR: Insertion of Organic Residual Products in culture systems (2018 – 2021)

Germany

24. Application-oriented carbon budget modelling of organic soils
25. BESTLAND-GHG reduction and soil Biodiversity in poorly drained soils under perennial crop
26. Soil hydraulic properties and release of CO₂ from peat soils
27. CarboCheck - A software tool to accumulate soil organic matter in agricultural soils
28. CarboHedge: Hedgerows and field copses in the emission inventories – Potential for carbon sequestration
29. Mitigating Agricultural Greenhouse Gas Emissions by improved pH management of soils (MAGGE-pH)
30. Eternal-C – Long term turnover and stabilisation of organic matter in topsoils and subsoils
31. VESBO - Impact assessment of vascular plant encroachment on water and carbon cycling in a Sphagnum dominated bog
32. National Contribution to the Integrated Carbon Observation System (ICOS)
33. Climate change mitigation through catch crops
34. Measuring and Modelling Greenhouse- Gas Emissions and nitrate leaching of raw material crop rotations (MASTER)
35. Effects of data requirements and uncertainty on prediction accuracy for model based evaluation and mitigation of nitrous oxide emissions from raw material cropping systems (THG-EMOBA).
36. Peatland monitoring program for climate protection
37. Reduction of NH₃ losses and improvement of nitrogen use efficiency after application of synthetic nitrogen fertilizers
38. SUBSOM - The Forgotten Part of Carbon Cycling: Organic Matter Storage and Turnover in Subsoils
39. Submerged drains in the model project "Gnarrenburger Moor"
40. Temporal trends of carbon stocks in agricultural soils
41. Water management for the reduction of GHG emissions from grasslands on peat soils (SWAMPS)
<https://www.thuenen.de/en/ak/projects/>
42. DESIRE project starting -Paludiculture in the Neman River
<https://www.moorwissen.de/en/index.php>



Hungary

43. Interdisciplinary Research Group for Promoting Climate-Smart and Sustainable Agriculture
44. 35. Evaluation of different management systems based on CO₂ and N₂O emissions
45. 36. Short-term and long-term responses of growth, carbon allocation and water balance of canopy tree species to fluctuation and extremes of climate conditions in mixed deciduous forest stands
46. 37. Effect of climate change on Hungarian humid, mesic and dry oak forests and their soils organic carbon storage capacity
47. Improving the greenhouse gas budget estimation for Hungary using tall tower observations
48. Measuring and modelling water and carbon balance of managed agricultural lands
49. Impact of spatial allocation of soil water and soil organic carbon on greenhouse gas emission in a small catchment (NKFIH - OTKA pályázat, témavezető: Horel Ágota)
50. Evaluation of different management systems based on CO₂ and N₂O emissions (NKFIH-OTKA pályázat, témavezető: Tóth Eszter)
51. Effect of soil management on soil carbon-dioxide emission at different Croatian and Hungarian agricultural sites" - (TÉT pályázat, témavezető: Tóth Eszter).
52. Compilation of indicator SO1 'Proportion of land that is degraded over total land area' from three sub-indicators for UNCCD reporting of Hungary
53. Methodology to establish an indicator of 'Trends in land productivity or functioning of the land' (SO1-2)
54. Elaboration of methods for calculation of indicator SO1-3 'Trends in carbon stocks above and below ground' for UNCCD reporting of Hungary
55. Quantification, evaluation and communication of uncertainty of predictive soil maps using advanced statistical approaches: On the example of soil organic carbon stock and its change.
56. Elaborating a methodology for estimating and mapping the potential organic carbon storage capacity of Hungarian soils

Ireland

57. Soil Quality Assessment & Research (SQUARE).
<https://www.teagasc.ie/environment/soil/research/square/>The focus of the SQUARE
58. project is to; further develop understanding of the soil functional capacity (incl. C sequestration) and quality concepts; assess the impact of soil structural degradation on the functional capacity of the soil.
59. The Agricultural Greenhouse Gas Research Initiative for Ireland (AGRI-I) is a consortium of researchers, students and professionals working collaboratively on multiple projects to develop verified strategies to decrease greenhouse gas emissions from Irish agriculture.
<https://agri-i.ie/projects/>
60. AGRI-SOC: Evaluating Land-Use and Land Management Impacts on Soil organic Carbon in Irish Agricultural Systems.

Italy

61. LIFE15 CCM/IT/000141 - OLIVE4CLIMATE - LIFE - OLIVE4CLIMATE - LIFE. CLIMATE CHANGE MITIGATION THROUGH A SUSTAINABLE SUPPLY CHAIN FOR THE OLIVE OIL SECTOR. Duration, 01/07/2016 – 30/12/2019: <https://olive4climate.eu/it/>
62. LIFE18 CCM/IT/001093 - LIFE agriCOLture - Livestock farming against climate change problems posed by soil degradation in the Emilian Apennines. Duration, 02/09/2019 – 31/08/2023:
https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPage&n_proj_id=7140
63. LIFE14 CCM/CY/000990 - LIFE+ ORGANIKO - Revamping organic farming and its products in



- the context of climate change mitigation strategies. Duration, 01/09/2015 – 31/08/2019:
64. https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPage&n_proj_id=5354
 65. LIFE15 PRE/IT/000001 - LIFE MEDINET - Mediterranean Network for Reporting Emissions and Removals in Cropland and Grassland. Duration, 01/04/2016 – 31/01/2018:
<https://www.lifemedinet.com/>
 66. LIFE17 CCM/GR/000087 - LIFE ClimaMed - Innovative technologies for climate change mitigation by Mediterranean agricultural sector. Duration, 01/07/2018 – 31/12/2022:
https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPage&n_proj_id=6814
 67. LIFE14 CCM/GR/000635 - CLIMATREE - A novel approach for accounting & monitoring carbon sequestration of tree crops and their potential as carbon sink areas. Duration: 16/07/2015 - 28/06/2020: <http://adapt2clima.eu/it/il-progetto>
 68. Horizon2020-DIVERFARMING. Duration, 01/05/2017 – 31/10/2022:
<http://www.diverfarming.eu/index.php/en/>
 69. Horizon2020-SoilCare - Soil Care for profitable and sustainable crop production in Europe. Duration, 01/03/2016 – 28/02/2021: <https://www.soilcare-project.eu/en/>
 70. Terras project Soil in sustainable agriculture - national project financed by Sardinia Region. Started in 2017: <http://www.sardegnaagricoltura.it/innovazionericerca/agris/>
 71. LIFE15 CCM/IT/000039 - LIFE+FORAGE4CLIMATE - FORAGE SYSTEMS FOR LESS GHG EMISSION AND MORE SOIL CARBON SINK IN CONTINENTAL AND MEDITERRANEAN AGRICULTURAL AREAS. Duration, 01/09/2016 – 31/08/2020:
http://forage4climate.crupa.it/nqcontent.cfm?a_id=14261
 72. LIFE17 CCM/IT/000062 - LIFE AGRESTIC - Reduction of Agricultural Greenhouse gases Emissions Through Innovative Cropping systems. Duration, 01/01/2019 – 30/06/2023:
<https://www.agrestic.eu/>
 73. LIFE + 12 ENV/IT/000578 HelpSoil. Duration, 01/07/2013 – 30/06/2017:
<http://www.lifehelpsoil.eu/>
 74. LIFE12 ENV/IT/000404 - LIFE+ Climate change R - Reduction of greenhouse gases from agricultural systems of Emilia-Romagna. Duration, 01/02/2010 – 31/12/2013:
https://ec.europa.eu/environment/life/project/Projects/index.cfm?fuseaction=search.dspPage&n_proj_id=4564 <https://progeu.regione.emilia-romagna.it/it/climatechanger>
 75. LIFE11 ENV/IT/000302 - IPNOA - Improved flux Prototypes for N2O emission reduction from Agriculture. Duration, 01/06/2012 – 30/11/2016: <http://www.ipnoa.eu/it/>
 76. 2014–2020 Rural Development Programme for Basilicata Region (Misura 16.1, PROSIT CUP H86G18000080002). Duration, 2017 – 2020:
<https://www.alsia.it/opencms/opencms/Temi/attivita/PRO.S.IT.-Vitivinicoltura/> (in Italian).

Latvia

77. [Climate change mitigation possibilities in organic soils](https://www.orgbalt.eu/?page_id=1719&lang=lv)
https://www.orgbalt.eu/?page_id=1719&lang=lv
78. Peat restoration project
<https://life-peat-restore.eu/lv/projekts/latvija/>

Lithuania

79. Investigation on development tendencies of soils with different genesis and geoecological potential under contrasting land management.



Netherlands

In the program Smart Soil Use projects have been started in 2018 to quantify the effect of C-sequestration in Dutch agricultural mineral soils:

80. Test field underwater drainage Friesland (Praktijkproeven onderwaterdrainage Friesland)
81. Pilot cattail in Marickenland (Pilot Iisdodde in Marickenland)
82. Calculating greenhousegas emissions for a waterarea plan Groot Wilnis-Vinkeveen (Berekening broeikasgasemissies voor watergebiedsplan Groot Wilnis-Vinkeveen)
83. National measurementcampaign greenhousegas emissions from peat (Landelijk meetprogramma broeikasgassen uit veen)
84. Fieldstudy effects of soil measures on peat oxidation (Veldonderzoek effecten bodemmaatregelen op veenoxidatie)

Norway

85. National project on the effects of reduced grassland management on SOC (ongoing, results by this/next year). Funded by the Norwegian Research Council:
<https://www.nibio.no/en/projects/carbon-storage-in-long-and-short-term-grasslands?locationfilter=true>
86. The PLATON project comprises a large platform in which agriculture and land uses are key elements. NIBIO participates in this project and will provide expertise with the aim to establish a plan for SOC monitoring in Norway as well as working towards the implementation of biochar as part of the GHG Inventory Report of Norway submitted to UNFCCC every year.: <https://www.platonklima.no/landbruk>

Poland

No information

Portugal

No information

Slovakia

87. Agricultural soil management with regard to its sustainability 2019-2021. Research project of the Ministry of Agriculture of Slovakia/2019-2021/Slovak: One of the important task of this project is modelling of SOC stock on another two agricultural farms. Project descriptions is not publically available
88. Scientific support of climate change adaptation in agriculture and mitigation of soil degradation (URANOS), Task A 3.2. Prediction of soil organic carbon stock in conditions of climate change. 1.4.2020 – 30.6.2023
89. Sustainable smart farming systems taking into account the future challenges (SMARTFARM), Activity 1: The optimization of cultivation systems from the viewpoint of production and non-production functions of the soil.1.10.2020 – 31.3.2023

Slovenia

No information

Spain

No information

Sweden



90. Many ongoing projects granted by FORMAS (a Swedish Research council focusing on management of natural resources) are addressing questions that are relevant in this context: <http://proj.formas.se/default.asp?funk=as>
91. Several more applied projects are granted by SLF (database only in Swedish): <http://www.lantbruksforskning.se/projektbanken/>
92. Several project granted by VR (Swedish research council for all more basic sciences) is also granting project related to the subject: <https://www.vr.se/english.html>
93. Carbon sequestration potential in top- and subsoil – analyzing soil databases and long-term trials.: http://www.lantbruksforskning.se/projektbanken/kolfastlaggningspotential-i-matjorden-och-alven-an/?app_year=&pub_year=&page=1&category=&search=k%C3%A4tterer
94. Turnover of organic matter in agricultural soil throughout the year: <http://proj.formas.se/detail.asp?arendeid=9272&x=250&y=20&sprak=1&redovisning=0>
95. Effects of soil organic carbon fractions on soil structure and preferential solute transport: <http://proj.formas.se/detail.asp?arendeid=9267&x=250&y=20&sprak=1&redovisning=0>
96. Mistra Food Futures: <https://www.mistra.org/nyhet/64-msek-till-forskning-om-hallbart-och-resilient-livsmedelssystem/>
97. Project PEATWISE: <https://www.eragas.eu/en/eragas/Research-projects/PEATWISE.htm>
98. KLIMASMART: www.slu.se/torv
99. MAGGEpH: <https://www.eragas.eu/en/eragas/Research-projects/MAGGE-pH.htm>
100. Fullständiga växthusgasbudgetar för odlade mulljordar skapar underlag för klimatsmarta åtgärder: http://www.lantbruksforskning.se/projektbanken/fullstandiga-vaxthusgasbudgetar-for-odlade-mulljor/?search=torv&app_year=&category=&page=1&pub_year=

Switzerland

101. Ongoing project on biochar addition to soil and its effect on SOC. This is not a classical research project, but rather a monitoring of farmer's practices on five fields.
102. Ongoing modelling study on C-sequestration potentials in Swiss agricultural soils considering crop rotation, biochar, agroforestry. The whole country will be covered, level of spatial disintegration is not yet decided on. Results are expected not before 2021.
103. Experiment on soil coverage: An ongoing project studies the GHG balance of an intensively managed organic soil under grassland with and without coverage with mineral soil. Results are expected in 2022.
104. Planned experiment on paddy rice on organic soil: A controlled experiment with organic soil in a mesocosm unit is planned to test these practices on their effect of GHG budgets: soil coverage, biochar addition, paddy rice with different water management.

Turkey

105. Soil Organic Carbon Project., TÜBİTAK BİLGEM, 2017-2018. (With the production of TOC stock map, gap analysis, areas without soil organic carbon data were identified and these areas were prioritized. In addition, Carbon Focused Biogeographical Regions were created within the scope of the Project).
106. Effects of Soil Tillage Techniques on Carbon Involvement and Sustainability of Soils. 2010-2017. Ministry of Agriculture and Forestry, DG of Agricultural Research and Policies (TAGEM), Soil, Fertilizer And Water Resources Central Research Institute (In this project; As a alternative to traditional tillage applications in fallow-wheat and leguminous-wheat crop rotation, the effects of reduced and zero tillage applications on increasing and maintaining soil efficiency and organic matter potential, have been tried to be put forward).
107. Determination of Seasonal and Annual Carbon Dioxide Output Amounts of Şanlıurfa Harran



- Plain Soils. 2017-2019. Ministry of Agriculture (TAGEM, GAP Agricultural Research Institute). (In this project; Amount of soil respiration was determined on different irrigated corn and cotton fields with tillaged and non-tillaged applications).
108. Turkey: Mapping Soil Carbon Stocks. 2016. Encyclopaedia of Soil Science, Third Edition DOI: 10.1081/E-ESS3-120052900
109. National Geospatial Soil Fertility and Soil Organic Carbon Information System Project (UTF/TUR/057/TUR) 2012-2015. Ministry of Agriculture and Forestry, DG of Agricultural Research and Polices(TAGEM), Soil, Fertilizer And Water Resources Central Research Institute-FAOSEC.

UK

110. UK Greenhouse Gas Inventory, 1990 to 2018: Annual Report for submission under the Framework Convention on Climate Change. https://uk-air.defra.gov.uk/assets/documents/reports/cat09/2004231028_ukghgi-90-18_Main_v02-00.pdf
111. NERC Project (NE/P019455/1): Soils Research to deliver Greenhouse Gas REmovals and Abatement Technologies (Soils-R-GGREAT) http://gotw.nerc.ac.uk/list_full.asp?pcode=NE%2FP019455%2F1
112. NERC Soil Security Programme (2015-2020) - a multi-million research investment to understand how soils resist, recover and ultimately adapt to land use and climate change. <https://soilsecurity.org/>
113. NERC Project (NE/S005137/1): 'LOCKED UP': The role of biotic and abiotic interactions in the stabilisation and persistence of soil organic carbon. http://gotw.nerc.ac.uk/list_full.asp?pcode=NE%2FS005137%2F1
114. The Soil Carbon project is an innovative project that aims to help farmers manage soils in a more sustainable and profitable way. <https://farmcarbontoolkit.org.uk/soil-carbon-project>



D List of experts and contacts Task T 2.4.3

No.	Participating organisation	Abbreviation	Country	Contact for T2 4.3
1	Institut National de la recherche Agronomique	INRAE	France	Sophie Cornu sophie.cornu@inrae.fr Sylvain Pellerin sylvain.pellerin@inrae.fr Eloïse Mason eloise.mason@inrae.fr
2	Wageningen Research	WR	The Netherlands	Peter Kuikman Peter.kuikman@wur.nl Thalisa Slier thalisa.slier@wur.nl Jan Peter Lesschen janpeter.lesschen@wur.nl
3	BIOS Science Austria	BIOS	Austria	Sophie Zechmeister Sophie.zechmeister@boku.ac.at
4	Flanders Research Institute for Agriculture, Fisheries and Food	EV-ILVO	Belgium	Maarten De Boever maarten.deboever@ilvo.vlaanderen.be
5	Centre Wallon de Recherches Agronomiques	CRAW	Belgium	Bruno Huyghebaert b.huyghebaert@cra.wallonie.be Brieuc Hardy b.hardy@cra.wallonie.be
6	Czech University of Life Sciences	CULS	Czech Republic	Lenka Pavlu pavlu@af.czu.cz
7	Aarhus University, Danish Centre for Food and Agriculture	AU	Denmark	Arezoo Taghizadeh arezoo.taghizadeh-toosi@agro.au.dk
8	Estonian University of Life Sciences	EMU	Estonia	Karin Kauer karin.kauer@emu.ee
9	Natural Resources Institute Finland	LUKE	Finland	Nils Kauer nils.borchard@luke.fi Markus Lier markus.lier@luke.fi
10	Johann Heinrich von Thünen-Institut	Theunen	Germany	Axel Don axel.don@thuenen.de Daria Seitz : daria.seitz@thuenen.de
12	Institute for Soil Sciences Centre for Agricultural Research (ATK TAKI)	ATK	Hungary	Toth Gergely toth.gergely@agrar.mta.hu Peter Laszlo laszlo.peter@agrar.mta.hu Zsafia Adrienn Kovacs kovacs.zsafia@atk.hu
13	Teagasc	Teagasc	Ireland	David Wall David.Wall@teagasc.ie, Lilian O'Sullivan Lilian.OSullivan@teagasc.ie
14	Council for Agricultural Research and Economics	CREA	Italy	Claudia Di Bene claudia.dibene@crea.gov.it
15	University of Latvia	UL	Latvia	Andis Lazdiņš andis.lazdins@silava.lv. Raimonds Kasparinskis raimonds.kasparinskis@lu.lv



16	Lithuanian Research Centre for Agriculture and Forestry	LAMMC	Lithuania	Dalia Feiziene dalia.feiziene@lammc.lt Kestutis Armolaitis kestutis.armolaitis@lammc.lt
17	Norwegian Institute of Bioeconomy Research	NIBIO	Norway	Teresa Gómez de la Bárcena Teresa.Barcelona@nibio.no Daniel Rasse daniel.rasse@nibio.no Alice Budai alice.budai@nibio.no
18	Institute of Soil Science and Plant Cultivation – State Research Institute	IUNG	Poland	Jacek Niedźwiecki: jacnk@iung.pulawy.pl Grzegorz Siebielec gs@iung.pulawy.pl
19	National Institute for Agrarian and Veterinarian Research I. P.	INIAV	Portugal	Corina Carranca corina.carranca@iniav.pt Raquel Mano raquel.mano@iniav.pt
20	National Agricultural and Food Centre	NPPC	Slovakia	Gabriela Barančíková gabriela.barancikova@nppc.sk
21	University of Ljubljana, Biotechnical Faculty, Centre for Soil and Environmental Science	ULBF	Slovenia	Rok Mihelič: rok.mihelic@bf.unilj.si
22	National Institute for Agriculture and Food Research and Technology	INIA	Spain	Elena Rodriguez rodriguez.elena@inia.es Benjamin Gimeno benjamins.gimeno@gmail.com
23	Swedish University of Agricultural Sciences	SLU	Sweden	Thomas Kätterer thomas.katterer@slu.se Bo Stenberg: Bo.Stenberg@slu.se
24	Agroscope	AGS	Switzerland	Jens Leifeld jens.leifeld@agroscope.admin.ch Leonor Rodrigues Leonormaria.gondimrodrigues@agroscope.admin.ch
25	Ministry of Agriculture and Forestry, General Directorate of Agricultural Research and Policies	TAGEM	Turkey	Sevinc Madenoglu: sevinc.madenoglu@tarimorman.gov.tr
26	Agri-Food and Biosciences Institute	AFBI	United	Dario Fornara dario.fornara@afbini.gov.uk



E Questionnaire for MS

Questionnaire Task 2.4.3

The **goal of T2.4.3.** is stocktaking and synthesis on available knowledge of **achievable carbon sequestration** in mineral soils, including pasture/grassland across Europe, under different farming systems, soil types and pedo-climatic conditions as well as on GHGs mitigation measures for managed organic soils under cropland or pasture/grassland.

The questionnaire below should give a framework for the kind of data and literature needed to response to the main objectives of the task. The information should include all kind of information, data, reports and literature on those subjects which are available **for each country** in any language. We kindly ask you to collect all the literature and data available and send it together with the completed questionnaire to Leonormaria.gondimrodrigues@agroscope.admin.ch and jens.leifeld@agroscope.admin.ch if possible before June.

Together with the data and questionnaire, please also deliver a summary in word format with a maximum of 3 pages excluding references, which include information on state-of-the-art knowledge on the achievable carbon sequestration in mineral soils in agricultural land, including pasture/grassland in your country, under different farming systems, soil types and pedo-climatic conditions as well as on GHGs mitigation measures for managed organic soils.

Also please indicate once again who is/are the appropriate contact persons and experts in your country (name, institution and contact details) for mineral soils and organic soils.

Achievable C-sequestration in MINERAL SOILS	
Here we are asking for the achievable carbon sequestration under a specific land management , and not for carbon stocks and soil carbon contents. C-sequestration as defined by Olson et al. (2014), “the process of transferring CO ₂ from the atmosphere into the soil of a land unit, through plants, plant residues and other organic solids which are stored or retained in the unit as part of the soil organic matter (humus) for a given time period.”	
MS country:	
EJP SOIL Partner:	
Experts and contact persons achievable C sequestration mineral soils:	
E-mail:	
Are there any quantitative estimations of the achievable carbon sequestration under different soil management in your country (finished or ongoing studies)?	
Yes	No
What is the form and accessibility of this information? (Published scientific or commercial literature, grey literature etc.)	Are there any studies planned for the future? Are there other studies comprising carbon storage potentials?
On which spatial scale did the studies operate? (Local, regional, national, environmental zones)	



Is this information available for the whole country or which regions have been studied?	
Please describe briefly, which methods were used to estimate or calculate the achievable carbon sequestration?	
If possible, please indicate which soil types have been considered for each study.	
For which period and duration has the C-sequestration been quantified?	
Are there any estimated C sequestration rates for different management practices derived from field studies in your country (finished or ongoing studies)?	
Yes	No
What is the form and accessibility of this information? (Published scientific or commercial literature, grey literature etc.)	Are there any other experiments or scenarios to estimate the achievable C-sequestration under different soil management using other C sequestration rates (e.g. global meta-analysis) relevant for your country?
What measures are involved?	
What are the potentials of these measures (e.g. rate/increase per year)?	
Potential for reducing GHG emissions from managed ORGANIC SOILS	
MS country:	
EJP SOIL Partner:	
Expert and contact person reducing GHG emissions from managed organic soils:	
E-mail:	
Are there any quantitative estimates for reducing GHG emissions (CO₂, CH₄, N₂O) from managed organic soils under different soil management in your country (finished or ongoing studies)?	
Yes	No
On which spatial scale did the studies operate? (Local, regional, national, environmental zones)	Are there other country-specific studies on mitigation of GHG emissions from managed organic soils in your country?
Is this information available for the whole country or which regions have been studied?	
Please describe briefly, which methods were used to estimate or calculate the GHG mitigation potential for managed organic soils.	
For which period and duration has the mitigation potential being quantified?	
Are there any estimated GHG mitigation potentials for managed organic soils derived from field studies in your country?	



Yes	No
What is the form and accessibility of this information? (Published scientific or commercial literature, grey literature etc.)	Are there any other experiments or scenarios to estimate the GHG mitigation potential for managed organic soils (e.g. global meta-analysis) relevant for your country?
What measures are involved?	
What are the potentials of these measures (e.g. rates per year)?	
Additional information	
What are the latest insights from your country over the last 5 years and which experiments and observations are running and results underway?	
Yes	No
Link to programmes and projects ongoing Do you have any project descriptions publically available?	
What are the most urgent knowledge gaps that you may have identified?	
Yes	No
Have these knowledge gaps been published and available to all?	

